

SYMMETRY



Claire Vallance
University of Oxford

Book: Symmetry (Vallance)

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (<https://LibreTexts.org>) and like the hundreds of other texts available within this powerful platform, it is freely available for reading, printing and "consuming." Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects.

Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



The LibreTexts mission is to unite students, faculty and scholars in a cooperative effort to develop an easy-to-use online platform for the construction, customization, and dissemination of OER content to reduce the burdens of unreasonable textbook costs to our students and society. The LibreTexts project is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning by developing an Open Access Resource environment. The project currently consists of 14 independently operating and interconnected libraries that are constantly being optimized by students, faculty, and outside experts to supplant conventional paper-based books. These free textbook alternatives are organized within a central environment that is both vertically (from advance to basic level) and horizontally (across different fields) integrated.

The LibreTexts libraries are Powered by [NICE CXOne](#) and are supported by the Department of Education Open Textbook Pilot Project, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptations contact info@LibreTexts.org. More information on our activities can be found via Facebook (<https://facebook.com/Libretexts>), Twitter (<https://twitter.com/libretexts>), or our blog (<http://Blog.Libretexts.org>).

This text was compiled on 01/10/2024

TABLE OF CONTENTS

Licensing

1: Chapters

- 1.1: Introduction to Symmetry
- 1.2: Symmetry Operations and Symmetry Elements
- 1.3: Symmetry Classification of Molecules- Point Groups
- 1.4: Symmetry and Physical Properties
- 1.5: Combining Symmetry Operations - 'Group Multiplication'
- 1.6: Constructing higher groups from simpler groups
- 1.7: Mathematical Definition of a Group
- 1.8: Review of Matrices
- 1.9: Transformation matrices
- 1.10: Matrix Representations of Groups
- 1.11: Properties of Matrix Representations
- 1.12: Reduction of Representations I
- 1.13: Irreducible representations and symmetry species
- 1.14: Character Tables
- 1.15: Reduction of representations II
- 1.16: Symmetry Adapted Linear Combinations (SALCs)
- 1.17: Determining whether an Integral can be Non-zero
- 1.18: Bonding in Diatomics
- 1.19: Bonding in Polyatomics- Constructing Molecular Orbitals from SALCs
- 1.20: Calculating Orbital Energies and Expansion Coefficients
- 1.21: Solving the Secular Equations
- 1.22: Summary of the Steps Involved in Constructing Molecular Orbitals
- 1.23: A more complicated bonding example
- 1.24: Molecular Vibrations
- 1.25: Summary of applying group theory to molecular motions
- 1.26: Group theory and Molecular Electronic States
- 1.27: Spectroscopy - Interaction of Atoms and Molecules with Light
- 1.28: Summary
- 1.29: Appendix A
- 1.30: Appendix B- Point Groups

Index

Glossary

Detailed Licensing

Licensing

A detailed breakdown of this resource's licensing can be found in [Back Matter/Detailed Licensing](#).

CHAPTER OVERVIEW

1: Chapters

- 1.1: Introduction to Symmetry
- 1.2: Symmetry Operations and Symmetry Elements
- 1.3: Symmetry Classification of Molecules- Point Groups
- 1.4: Symmetry and Physical Properties
- 1.5: Combining Symmetry Operations - 'Group Multiplication'
- 1.6: Constructing higher groups from simpler groups
- 1.7: Mathematical Definition of a Group
- 1.8: Review of Matrices
- 1.9: Transformation matrices
- 1.10: Matrix Representations of Groups
- 1.11: Properties of Matrix Representations
- 1.12: Reduction of Representations I
- 1.13: Irreducible representations and symmetry species
- 1.14: Character Tables
- 1.15: Reduction of representations II
- 1.16: Symmetry Adapted Linear Combinations (SALCs)
- 1.17: Determining whether an Integral can be Non-zero
- 1.18: Bonding in Diatomics
- 1.19: Bonding in Polyatomics- Constructing Molecular Orbitals from SALCs
- 1.20: Calculating Orbital Energies and Expansion Coefficients
- 1.21: Solving the Secular Equations
- 1.22: Summary of the Steps Involved in Constructing Molecular Orbitals
- 1.23: A more complicated bonding example
- 1.24: Molecular Vibrations
- 1.25: Summary of applying group theory to molecular motions
- 1.26: Group theory and Molecular Electronic States
- 1.27: Spectroscopy - Interaction of Atoms and Molecules with Light
- 1.28: Summary
- 1.29: Appendix A
- 1.30: Appendix B- Point Groups

This page titled [1: Chapters](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.1: Introduction to Symmetry

You will already be familiar with the concept of symmetry in an everyday sense. If we say something is ‘symmetrical’, we usually mean it has mirror symmetry, or ‘left-right’ symmetry, and would look the same if viewed in a mirror. Symmetry is also very important in chemistry. Some molecules are clearly ‘more symmetrical’ than others, but what consequences does this have, if any?

The aim of this course is to provide a systematic treatment of symmetry in chemical systems within the mathematical framework known as *group theory* (the reason for the name will become apparent later on). Once we have classified the symmetry of a molecule, group theory provides a powerful set of tools that provide us with considerable insight into many of its chemical and physical properties.

Some applications of group theory that will be covered in this course include:


1. Predicting whether a given molecule will be chiral, or polar.
2. Examining chemical bonding and visualising molecular orbitals.
3. Predicting whether a molecule may absorb light of a given polarisation, and which spectroscopic transitions may be excited if it does.
4. Investigating the vibrational motions of the molecule.

You may well meet some of these topics again, possibly in more detail, in later courses. However, they will be introduced here to give you a fairly broad introduction to the capabilities and applications of group theory once we have worked through the basic principles and ‘machinery’ of the theory.

This page titled [1.1: Introduction to Symmetry](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.2: Symmetry Operations and Symmetry Elements


A *symmetry operation* is an action that leaves an object looking the same after it has been carried out. For example, if we take a molecule of water and rotate it by 180° about an axis passing through the central O atom (between the two H atoms) it will look the same as before. It will also look the same if we reflect it through either of two mirror planes, as shown in the figure below.



Each symmetry operation has a corresponding *symmetry element*, which is the axis, plane, line or point with respect to which the symmetry operation is carried out. The symmetry element consists of all the points that stay in the same place when the symmetry operation is performed. In a rotation, the line of points that stay in the same place constitute a *symmetry axis*; in a reflection the points that remain unchanged make up a *plane of symmetry*.

The symmetry elements that a molecule may possess are:

1. E - the identity. The identity operation consists of doing nothing, and the corresponding symmetry element is the entire molecule. Every molecule has at least this element.
2. C_n - an n -fold axis of rotation. Rotation by $360^\circ/n$ leaves the molecule unchanged. The H_2O molecule above has a C_2 axis. Some molecules have more than one C_n axis, in which case the one with the highest value of n is called the *principal axis*. Note that by convention rotations are *counterclockwise* about the axis.
3. σ - a plane of symmetry. Reflection in the plane leaves the molecule looking the same. In a molecule that also has an axis of symmetry, a mirror plane that includes the axis is called a vertical mirror plane and is labeled σ_v , while one perpendicular to the axis is called a horizontal mirror plane and is labeled σ_h . A vertical mirror plane that bisects the angle between two C_2 axes is called a dihedral mirror plane, σ_d .
4. i - a center of symmetry. Inversion through the center of symmetry leaves the molecule unchanged. Inversion consists of passing each point through the center of inversion and out to the same distance on the other side of the molecule. An example of a molecule with a center of inversion is shown below.



5. S_n - an n -fold improper rotation axis (also called a rotary-reflection axis). The rotary reflection operation consists of rotating through an angle $360^\circ/n$ about the axis, followed by reflecting in a plane perpendicular to the axis. Note that S_1 is the same as reflection and S_2 is the same as inversion. The molecule shown above has two S_2 axes.

The identity E and rotations C_n are symmetry operations that could actually be carried out on a molecule. For this reason they are called *proper symmetry operations*. Reflections, inversions and improper rotations can only be imagined (it is not actually possible to turn a molecule into its mirror image or to invert it without some fairly drastic rearrangement of chemical bonds) and as such, are termed *improper symmetry operations*.

Axis Definitions

Conventionally, when imposing a set of Cartesian axes on a molecule (as we will need to do later on in the course), the z axis lies along the principal axis of the molecule, the x axis lies in the plane of the molecule (or in a plane containing the largest number of atoms if the molecule is non-planar), and the y axis makes up a right handed axis system.

This page titled [1.2: Symmetry Operations and Symmetry Elements](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.


1.3: Symmetry Classification of Molecules- Point Groups

It is only possible for certain combinations of symmetry elements to be present in a molecule (or any other object). As a result, we may group together molecules that possess the same symmetry elements and classify molecules according to their symmetry. These groups of symmetry elements are called *point groups* (due to the fact that there is at least one point in space that remains unchanged no matter which symmetry operation from the group is applied). There are two systems of notation for labeling symmetry groups, called the Schoenflies and Hermann-Mauguin (or International) systems. The symmetry of individual molecules is usually described using the Schoenflies notation, and we shall be using this notation for the remainder of the course¹.


*Some of the **point groups** share their names with symmetry **operations**, so be careful you do not mix up the two. It is usually clear from the context which one is being referred to.*

Molecular Point Groups


1. C_1 - contains only the identity (a C_1 rotation is a rotation by 360° and is the same as the identity operation E) e.g. CHDFCl.



2. C_i - contains the identity E and a center of inversion i .



3. C_s - contains the identity E and a plane of reflection σ .




4. C_n - contains the identity and an n -fold axis of rotation.



5. C_{nv} - contains the identity, an n -fold axis of rotation, and n vertical mirror planes σ_v .



6. C_{nh} - contains the identity, an n -fold axis of rotation, and a horizontal reflection plane σ_h (note that in C_{2h} this combination of symmetry elements automatically implies a center of inversion).




7. D_n - contains the identity, an n -fold axis of rotation, and n 2-fold rotations about axes perpendicular to the principal axis.



8. D_{nh} - contains the same symmetry elements as D_n with the addition of a horizontal mirror plane.




9. D_{nd} - contains the same symmetry elements as D_n with the addition of n dihedral mirror planes.



10. S_n - contains the identity and *one* S_n axis. Note that molecules only belong to S_n if they have not already been classified in terms of one of the preceding point groups (e.g. S_2 is the same as C_i , and a molecule with this symmetry would already have been classified).

The following groups are the cubic groups, which contain more than one principal axis. They separate into the tetrahedral groups (T_d , T_h and T) and the octahedral groups (O and O_h). The icosahedral group also exists, but is not included below.

11. T_d - contains all the symmetry elements of a regular tetrahedron, including the identity, 4 C_3 axes, 3 C_2 axes, 6 dihedral mirror planes, and 3 S_4 axes e.g. CH_4 .




12. T - as for T_d but no planes of reflection.
 13. T_h - as for T but contains a center of inversion.
 14. O_h - the group of the regular octahedron e.g. SF_6 .



15. O - as for O_h , but with no planes of reflection.

The final group is the full rotation group R_3 , which consists of an infinite number of C_n axes with all possible values of n and describes the symmetry of a sphere. Atoms (but no molecules) belong to R_3 , and the group has important applications in atomic quantum mechanics. However, we won't be treating it any further here.

Once you become more familiar with the symmetry elements and point groups described above, you will find it quite straightforward to classify a molecule in terms of its point group. In the meantime, the flowchart shown below provides a step-by-step approach to the problem.



¹Though the Hermann-Mauguin system can be used to label point groups, it is usually used in the discussion of crystal symmetry. In crystals, in addition to the symmetry elements described above, translational symmetry elements are very important. Translational symmetry operations leave no point unchanged, with the consequence that crystal symmetry is described in terms of *space groups* rather than *point groups*.

This page titled [1.3: Symmetry Classification of Molecules- Point Groups](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.4: Symmetry and Physical Properties

Carrying out a symmetry operation on a molecule must not change any of its physical properties. It turns out that this has some interesting consequences, allowing us to predict whether or not a molecule may be chiral or polar on the basis of its point group.

Polarity

For a molecule to have a permanent dipole moment, it must have an asymmetric charge distribution. The point group of the molecule not only determines whether the molecule may have a dipole moment, but also in which direction(s) it may point.

If a molecule has a C_n axis with $n > 1$, it **cannot** have a dipole moment perpendicular to the axis of rotation (for example, a C_2 rotation would interchange the ends of such a dipole moment and reverse the polarity, which is not allowed – rotations with higher values of n would also change the direction in which the dipole points). Any dipole must lie parallel to a C_n axis.

Also, if the point group of the molecule contains any symmetry operation that would interchange the two ends of the molecule, such as a σ_h mirror plane or a C_2 rotation perpendicular to the principal axis, then there cannot be a dipole moment along the axis.

The only groups compatible with a dipole moment are C_n , C_{nv} and C_s . In molecules belonging to C_n or C_{nv} the dipole must lie along the axis of rotation.

Chirality

One example of symmetry in chemistry that you will already have come across is found in the isomeric pairs of molecules called enantiomers. Enantiomers are non-superimposable mirror images of each other, and one consequence of this symmetrical relationship is that they rotate the plane of polarized light passing through them in opposite directions. Such molecules are said to be chiral,² meaning that they cannot be superimposed on their mirror image. Formally, the symmetry element that precludes a molecule from being chiral is a rotation-reflection axis S_n . Such an axis is often implied by other symmetry elements present in a group.


For example, a point group that has C_n and σ_h as elements will also have S_n . Similarly, a center of inversion is equivalent to S_2 . As a rule of thumb, a molecule definitely cannot be chiral if it has a center of inversion or a mirror plane of any type (σ_h , σ_v or σ_d), but if these symmetry elements are absent the molecule should be checked carefully for an S_n axis before it is assumed to be chiral.

² The word *chiral* has its origins in the Greek word for hand ($\chi\epsilon\rho\iota$, pronounced 'cheri' with a soft ch as in 'loch'). A pair of hands is also a pair of non-superimposable mirror images, and you will often hear chirality referred to as 'handedness' for this reason.


This page titled [1.4: Symmetry and Physical Properties](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.5: Combining Symmetry Operations - 'Group Multiplication'

Now we will investigate what happens when we apply two symmetry operations in sequence. As an example, consider the NH_3 molecule, which belongs to the C_{3v} point group. Consider what happens if we apply a C_3 rotation followed by a σ_v reflection. We write this combined operation $\sigma_v C_3$ (when written, symmetry operations operate on the thing directly to their right, just as operators do in quantum mechanics – we therefore have to work backwards from right to left from the notation to get the correct order in which the operations are applied). As we shall soon see, the order in which the operations are applied is important.



The combined operation $\sigma_v C_3$ is equivalent to σ_v'' , which is also a symmetry operation of the C_{3v} point group. Now let's see what happens if we apply the operators in the reverse order i.e. $C_3 \sigma_v$ (σ_v followed by C_3).



Again, the combined operation $C_3 \sigma_v$ is equivalent to another operation of the point group, this time σ_v' .

There are two important points that are illustrated by this example:

1. The order in which two operations are applied is important. For two symmetry operations A and B , AB is not necessarily the same as BA , i.e. symmetry operations do not in general commute. In some groups the symmetry elements do commute; such groups are said to be Abelian.
2. If two operations from the same point group are applied in sequence, the result will be equivalent to another operation from the point group. Symmetry operations that are related to each other by other symmetry operations of the group are said to belong to the same *class*. In NH_3 , the three mirror planes σ_v , σ_v' and σ_v'' belong to the same class (related to each other through a C_3 rotation), as do the rotations C_3^+ and C_3^- (anticlockwise and clockwise rotations about the principal axis, related to each other by a vertical mirror plane).

The effects of applying two symmetry operations in sequence within a given point group are summarized in *group multiplication tables*. As an example, the complete group multiplication table for C_{3v} using the symmetry operations as defined in the figures above is shown below. The operations written along the first row of the table are carried out first, followed by those written in the first column (note that the table would change if we chose to name σ_v , σ_v' and σ_v'' in some different order).

C_{3v}	E	C_3^+	C_3^-	σ_v	σ_v'	σ_v''
E	E	C_3^+	C_3^-	σ_v	σ_v'	σ_v''
C_3^+	C_3^+	C_3^-	E	σ_v''	σ_v'	σ_v
C_3^-	C_3^-	E	C_3^+	σ_v'	σ_v	σ_v''
σ_v	σ_v	σ_v''	σ_v'	E	C_3^-	C_3^+
σ_v'	σ_v'	σ_v	σ_v''	C_3^+	E	C_3^-
σ_v''	σ_v''	σ_v'	σ_v	C_3^-	C_3^+	E

(1.5.1)

This page titled [1.5: Combining Symmetry Operations - 'Group Multiplication'](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.6: Constructing higher groups from simpler groups

A group that contains a large number of symmetry elements may often be constructed from simpler groups. This is probably best illustrated using an example. Consider the point groups C_2 and C_S . C_2 contains the elements E and C_2 , and has order 2, while C_S contains E and σ and also has order 2. We can use these two groups to construct the group C_{2v} by applying the symmetry operations of C_2 and C_S in sequence.

$$\begin{array}{llll} C_2 \text{ operation} & E & E & C_2 & C_2 \\ C_S \text{ operation} & E & \sigma(xz) & E & \sigma(xz) \\ \text{Result} & E & \sigma_v(xz) & C_2 & \sigma'_v(yz) \end{array} \quad (6.1)$$

Notice that C_{2v} has order 4, which is the product of the orders of the two lower-order groups. C_{2v} may be described as a direct product group of C_2 and C_S . The origin of this name should become obvious when we review the properties of matrices.

This page titled [1.6: Constructing higher groups from simpler groups](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.7: Mathematical Definition of a Group

Now that we have explored some of the properties of symmetry operations and elements and their behavior within point groups, we are ready to introduce the formal mathematical definition of a group.

A mathematical group is defined as a set of elements ($g_1, g_2, g_3 \dots$) together with a rule for forming combinations $g_i g_j$. The number of elements h is called the *order* of the group. For our purposes, the elements are the symmetry operations of a molecule and the rule for combining them is the sequential application of symmetry operations investigated in the previous section. The elements of the group and the rule for combining them must satisfy the following criteria.

1. The group must include the identity E , for which

$$Eg_i = g_i \quad (7.1)$$

for all the elements of the group.

2. The elements must satisfy the *group property* that the combination of any pair of elements is also an element of the group.
3. Each element g_i must have an inverse g_i^{-1} , which is also an element of the group, such that

$$g_i g_i^{-1} = g_i^{-1} g_i = E \quad (7.2)$$

(e.g. in C_{3v} the inverse of C_3^+ is C_3^- , the inverse of (σ_v is σ_v), the inverse g_i^{-1} effectively 'undoes' the effect of the symmetry operation g_i).

4. The rule of combination must be associative i.e.

$$(g_i g_j)(g_k) = g_i(g_j g_k) \quad (7.3)$$

The above definition **does not** require the elements to commute, which would require

$$g_i g_k = g_k g_i \quad (7.4)$$

As we discovered in the C_{3v} example above, in many groups the outcome of consecutive application of two symmetry operations depends on the order in which the operations are applied. Groups for which the elements do not commute are called *non-Abelian* groups; those for which they elements do commute are *Abelian*.

Group theory is an important area in mathematics, and luckily for chemists the mathematicians have already done most of the work for us. Along with the formal definition of a group comes a comprehensive mathematical framework that allows us to carry out a rigorous treatment of symmetry in molecular systems and learn about its consequences.

Many problems involving operators or operations (such as those found in quantum mechanics or group theory) may be reformulated in terms of matrices. Any of you who have come across transformation matrices before will know that symmetry operations such as rotations and reflections may be represented by matrices. It turns out that the set of matrices representing the symmetry operations in a group obey all the conditions laid out above in the mathematical definition of a group, and using matrix representations of symmetry operations simplifies carrying out calculations in group theory. Before we learn how to use matrices in group theory, it will probably be helpful to review some basic definitions and properties of matrices.

This page titled [1.7: Mathematical Definition of a Group](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.8: Review of Matrices

Definitions

An $n \times m$ matrix is a two dimensional array of numbers with n rows and m columns. The integers n and m are called the *dimensions* of the matrix. If $n = m$ then the matrix is *square*. The numbers in the matrix are known as *matrix elements* (or just elements) and are usually given subscripts to signify their position in the matrix e.g. an element a_{ij} would occupy the i^{th} row and j^{th} column of the matrix. For example:

$$M = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \quad (1.8.1)$$

is a 3×3 matrix with $a_{11} = 1$, $a_{12} = 2$, $a_{13} = 3$, $a_{21} = 4$ etc

In a square matrix, *diagonal* elements are those for which $i=j$ (the numbers 1, 5, and 9 in the above example). *Off-diagonal* elements are those for which $i \neq j$ (2, 3, 4, 6, 7, and 8 in the above example). If all the off-diagonal elements are equal to zero then we have a *diagonal* matrix. We will see later that diagonal matrices are of considerable importance in group theory.

A *unit matrix* or *identity matrix* (usually given the symbol I) is a diagonal matrix in which all the diagonal elements are equal to 1. A unit matrix acting on another matrix has no effect – it is the same as the identity operation in group theory and is analogous to multiplying a number by 1 in everyday arithmetic.

The *transpose* A^T of a matrix A is the matrix that results from interchanging all the rows and columns. A *symmetric* matrix is the same as its transpose ($A^T = A$ i.e. $a_{ij} = a_{ji}$ for all values of i and j). The transpose of matrix M above (which is not symmetric) is

$$M^T = \begin{pmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{pmatrix} \quad (1.8.2)$$

The sum of the diagonal elements in a square matrix is called the *trace* (or *character*) of the matrix (for the above matrix, the trace is $\chi = 1 + 5 + 9 = 15$). The traces of matrices representing symmetry operations will turn out to be of great importance in group theory.

A *vector* is just a special case of a matrix in which one of the dimensions is equal to 1. An $n \times 1$ matrix is a *column vector*; a $1 \times m$ matrix is a *row vector*. The components of a vector are usually only labeled with one index. A *unit vector* has one element equal to 1 and the others equal to zero (it is the same as one row or column of an identity matrix). We can extend the idea further to say that a single number is a matrix (or vector) of dimension 1×1 .

Matrix Algebra

- i. Two matrices with the same dimensions may be added or subtracted by adding or subtracting the elements occupying the same position in each matrix. e.g.

$$A = \begin{pmatrix} 1 & 0 & 2 \\ 2 & 2 & 1 \\ 3 & 2 & 0 \end{pmatrix} \quad (1.8.3)$$

$$B = \begin{pmatrix} 2 & 0 & -2 \\ 1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix} \quad (1.8.4)$$

$$A + B = \begin{pmatrix} 3 & 0 & 0 \\ 3 & 2 & 2 \\ 4 & 1 & 0 \end{pmatrix} \quad (1.8.5)$$

$$A - B = \begin{pmatrix} -1 & 0 & 4 \\ 1 & 2 & 0 \\ 2 & 3 & 0 \end{pmatrix} \quad (1.8.6)$$

ii. A matrix may be multiplied by a constant by multiplying each element by the constant.

$$4B = \begin{pmatrix} 8 & 0 & -8 \\ 4 & 0 & 4 \\ 4 & -4 & 0 \end{pmatrix} \quad (1.8.7)$$

$$3A = \begin{pmatrix} 3 & 0 & 6 \\ 6 & 6 & 3 \\ 9 & 6 & 0 \end{pmatrix} \quad (1.8.8)$$

iii. Two matrices may be multiplied together provided that the number of columns of the first matrix is the same as the number of rows of the second matrix i.e. an $n \times m$ matrix may be multiplied by an $m \times l$ matrix. The resulting matrix will have dimensions $n \times l$. To find the element a_{ij} in the product matrix, we take the dot product of row i of the first matrix and column j of the second matrix (i.e. we multiply consecutive elements together from row i of the first matrix and column j of the second matrix and add them together i.e. $c_{ij} = \sum_k a_{ik}b_{jk}$ e.g. in the 3×3 matrices A and B used in the above examples, the first element in the product matrix $C = AB$ is $c_{11} = a_{11}b_{11} + a_{12}b_{21} + a_{13}b_{31}$

$$AB = \begin{pmatrix} 1 & 0 & 2 \\ 2 & 2 & 1 \\ 3 & 2 & 0 \end{pmatrix} \begin{pmatrix} 2 & 0 & -2 \\ 1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix} = \begin{pmatrix} 4 & -2 & -2 \\ 7 & -1 & -2 \\ 8 & 0 & -4 \end{pmatrix} \quad (1.8.9)$$

An example of a matrix multiplying a vector is

$$A\mathbf{v} = \begin{pmatrix} 1 & 0 & 2 \\ 2 & 2 & 1 \\ 3 & 2 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 7 \\ 9 \\ 7 \end{pmatrix} \quad (1.8.10)$$

Matrix multiplication is not generally commutative, a property that mirrors the behavior found earlier for symmetry operations within a point group.

Direct Products

The direct product of two matrices (given the symbol \otimes) is a special type of matrix product that generates a matrix of higher dimensionality if both matrices have dimension greater than one. The easiest way to demonstrate how to construct a direct product of two matrices A and B is by an example:

$$A \otimes B = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \otimes \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \quad (1.8.11)$$

$$= \begin{pmatrix} a_{11}B & a_{12}B \\ a_{21}B & a_{22}B \end{pmatrix} \quad (1.8.12)$$

$$= \begin{pmatrix} a_{11}b_{11} & a_{11}b_{12} & a_{12}b_{11} & a_{12}b_{12} \\ a_{11}b_{21} & a_{11}b_{22} & a_{12}b_{21} & a_{12}b_{22} \\ a_{21}b_{11} & a_{21}b_{12} & a_{22}b_{11} & a_{22}b_{12} \\ a_{21}b_{21} & a_{21}b_{22} & a_{22}b_{21} & a_{22}b_{22} \end{pmatrix} \quad (1.8.13)$$

Though this may seem like a somewhat strange operation to carry out, direct products crop up a great deal in group theory.


Inverse Matrices and Determinants

If two square matrices A and B multiply together to give the identity matrix I (i.e. $AB = I$) then B is said to be the *inverse* of A (written A^{-1}). If B is the inverse of A then A is also the inverse of B . Recall that one of the conditions imposed upon the symmetry operations in a group is that each operation must have an inverse. It follows by analogy that any matrices we use to represent symmetry elements must also have inverses. It turns out that a square matrix only has an inverse if its determinant is non-

zero. For this reason (and others which will become apparent later on when we need to solve equations involving matrices) we need to learn a little about matrix determinants and their properties.

For every square matrix, there is a unique function of all the elements that yields a single number called the determinant. Initially it probably won't be particularly obvious why this number should be useful, but matrix determinants are of great importance both in pure mathematics and in a number of areas of science. Historically, determinants were actually around before matrices. They arose originally as a property of a system of linear equations that 'determined' whether the system had a unique solution. As we shall see later, when such a system of equations is recast as a matrix equation this property carries over into the matrix determinant.

There are two different definitions of a determinant, one geometric and one algebraic. In the geometric interpretation, we consider the numbers across each row of an $n \times n$ matrix as coordinates in n -dimensional space. In a one-dimensional matrix (i.e. a number), there is only one coordinate, and the determinant can be interpreted as the (signed) length of a vector from the origin to this point. For a 2×2 matrix we have two coordinates in a plane, and the determinant is the (signed) area of the parallelogram that includes these two points and the origin. For a 3×3 matrix the determinant is the (signed) volume of the parallelepiped that includes the three points (in three-dimensional space) defined by the matrix and the origin. This is illustrated below. The idea extends up to higher dimensions in a similar way. In some sense then, the determinant is therefore related to the size of a matrix.



The algebraic definition of the determinant of an $n \times n$ matrix is a sum over all the possible products (permutations) of n elements taken from different rows and columns. The number of terms in the sum is $n!$, the number of possible permutations of n values (i.e. 2 for a 2×2 matrix, 6 for a 3×3 matrix etc). Each term in the sum is given a positive or a negative sign depending on whether the number of *permutation inversions* in the product is even or odd. A permutation inversion is just a pair of elements that are out of order when described by their indices. For example, for a set of four elements (a_1, a_2, a_3, a_4) , the permutation $a_1 a_2 a_3 a_4$ has all the elements in their correct order (i.e. in order of increasing index). However, the permutation $a_2 a_4 a_1 a_3$ contains the permutation inversions $a_2 a_1, a_4 a_1, a_4 a_3$.

For example, for a two-dimensional matrix

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \quad (1.8.14)$$

where the subscripts label the row and column positions of the elements, there are 2 possible products/permutations involving elements from different rows and column, $a_{11}a_{22}$ and $a_{12}a_{21}$. In the second term, there is a permutation inversion involving the column indices 2 and 1 (permutation inversions involving the row and column indices should be looked for separately) so this term takes a negative sign, and the determinant is $a_{11}a_{22} - a_{12}a_{21}$.

For a 3×3 matrix

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \quad (1.8.15)$$

the possible combinations of elements from different rows and columns, together with the sign from the number of permutations required to put their indices in numerical order are:

$$\begin{aligned} & a_{11}a_{22}a_{33} \quad (0 \text{ inversions}) \\ & -a_{11}a_{23}a_{32} \quad (1 \text{ inversion} - 3 > 2 \text{ in the column indices}) \\ & -a_{12}a_{21}a_{33} \quad (1 \text{ inversion} - 2 > 1 \text{ in the column indices}) \\ & a_{12}a_{23}a_{31} \quad (2 \text{ inversions} - 2 > 1 \text{ and } 3 > 1 \text{ in the column indices}) \\ & a_{13}a_{21}a_{32} \quad (2 \text{ inversions} - 3 > 1 \text{ and } 3 > 2 \text{ in the column indices}) \\ & -a_{13}a_{22}a_{31} \quad (3 \text{ inversions} - 3 > 2, 3 > 1, \text{ and } 2 > 1 \text{ in the column indices}) \end{aligned} \quad (1.8.16)$$

and the determinant is simply the sum of these terms.

This may all seem a little complicated, but in practice there is a fairly systematic procedure for calculating determinants. The determinant of a matrix A is usually written $\det(A)$ or $|A|$.

For a 2×2 matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}; \det(A) = |A| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc \quad (1.8.17)$$

For a 3×3 matrix

$$B = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}; \det(B) = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix} \quad (1.8.18)$$

For a 4×4 matrix

$$C = \begin{pmatrix} a & b & c & d \\ e & f & g & h \\ i & j & k & l \\ m & n & o & p \end{pmatrix}; \det(C) = a \begin{vmatrix} f & g & h \\ j & k & l \\ n & o & p \end{vmatrix} - b \begin{vmatrix} e & g & h \\ i & k & l \\ m & o & p \end{vmatrix} + c \begin{vmatrix} e & f & h \\ i & j & l \\ m & n & p \end{vmatrix} - d \begin{vmatrix} e & f & g \\ i & j & k \\ m & n & o \end{vmatrix} \quad (1.8.19)$$

and so on in higher dimensions. Note that the submatrices in the 3×3 example above are just the matrices formed from the original matrix B that don't include any elements from the same row or column as the premultiplying factors from the first row.

Matrix determinants have a number of important properties:

i. The determinant of the identity matrix is 1.

$$e.g. \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 1 \quad (1.8.20)$$

ii. The determinant of a matrix is the same as the determinant of its transpose i.e. $\det(a) = \det(A^T)$

$$e.g. \begin{vmatrix} a & b \\ c & d \end{vmatrix} = \begin{vmatrix} a & c \\ b & d \end{vmatrix} \quad (1.8.21)$$

iii. The determinant changes sign when any two rows or any two columns are interchanged

$$e.g. \begin{vmatrix} a & b \\ c & d \end{vmatrix} = - \begin{vmatrix} b & a \\ d & c \end{vmatrix} = - \begin{vmatrix} c & d \\ a & b \end{vmatrix} = \begin{vmatrix} d & c \\ b & a \end{vmatrix} \quad (1.8.22)$$

iv. The determinant is zero if any row or column is entirely zero, or if any two rows or columns are equal or a multiple of one another.

$$e.g. \begin{vmatrix} 1 & 2 \\ 0 & 0 \end{vmatrix} = 0, \begin{vmatrix} 1 & 2 \\ 2 & 4 \end{vmatrix} = 0 \quad (1.8.23)$$

v. The determinant is unchanged by adding any linear combination of rows (or columns) to another row (or column).

vi. The determinant of the product of two matrices is the same as the product of the determinants of the two matrices i.e. $\det(AB) = \det(A)\det(B)$.

The requirement that in order for a matrix to have an inverse it must have a non-zero determinant follows from property vi). As mentioned previously, the product of a matrix and its inverse yields the identity matrix I . We therefore have:

$$\begin{aligned} \det(A^{-1}A) &= \det(A^{-1})\det(A) = \det(I) \\ \det(A^{-1}) &= \det(I)/\det(A) = 1/\det(A) \end{aligned} \quad (1.8.24)$$

It follows that a matrix A can only have an inverse if its determinant is non-zero, otherwise the determinant of its inverse would be undefined.

This page titled [1.8: Review of Matrices](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.9: Transformation matrices

Matrices can be used to map one set of coordinates or functions onto another set. Matrices used for this purpose are called *transformation matrices*. In group theory, we can use transformation matrices to carry out the various symmetry operations considered at the beginning of the course. As a simple example, we will investigate the matrices we would use to carry out some of these symmetry operations on a vector (x, y) .

The identity Operation

The identity operation leaves the vector unchanged, and as you may already suspect, the appropriate matrix is the identity matrix.

$$(x, y) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = (x, y) \quad (1.9.1)$$

Reflection in a plane

The simplest example of a reflection matrix corresponds to reflecting the vector (x, y) in either the x or y axes. Reflection in the x axis maps y to $-y$, while reflection in the y axis maps x to $-x$. The appropriate matrix is very like the identity matrix but with a change in sign for the appropriate element. Reflection in the x axis transforms the vector (x, y) to $(x, -y)$, and the appropriate matrix is

$$(x, y) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = (x, -y) \quad (1.9.2)$$




Figure 1.9.1: Reflection across the x-axis

Reflection in the y axis transforms the vector (x, y) to $(-x, y)$, and the appropriate matrix is

$$(x, y) \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = (-x, y) \quad (1.9.3)$$




Figure 1.9.2: Reflection across the y-axis

More generally, matrices can be used to represent reflections in any plane (or line in 2D). For example, reflection in the 45° axis shown below maps

(x, y) onto $(-y, -x)$.

$$(x, y) \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} = (-y, -x) \quad (1.9.4)$$




Figure 1.9.3: Reflection across the axis that is rotated 45° with respect to x-axis.

Rotation about an Axis

In two dimensions, the appropriate matrix to represent rotation by an angle θ about the origin is

$$R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \quad (1.9.5)$$

In three dimensions, rotations about the x , y and z axes acting on a vector (x, y, z) are represented by the following matrices.

$$R_x(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\theta & -\sin\theta \\ 0 & \sin\theta & \cos\theta \end{pmatrix} \quad (1.9.6)$$

$$R_y(\theta) = \begin{pmatrix} \cos\theta & 0 & -\sin\theta \\ 0 & 1 & 0 \\ \sin\theta & 0 & \cos\theta \end{pmatrix} \quad (1.9.7)$$

$$R_z(\theta) = \begin{pmatrix} \cos\theta & -\sin\theta & 0 \\ \sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (1.9.8)$$

This page titled [1.9: Transformation matrices](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.10: Matrix Representations of Groups


We are now ready to integrate what we have just learned about matrices with group theory. The symmetry operations in a group may be represented by a set of transformation matrices $\Gamma(g)$, one for each symmetry element g . Each individual matrix is called a **representative** of the corresponding symmetry operation, and the complete set of matrices is called a **matrix representation** of the group. The matrix representatives act on some chosen **basis set** of functions, and the actual matrices making up a given representation will depend on the basis that has been chosen. The representation is then said to **span** the chosen basis. In the examples above we were looking at the effect of some simple transformation matrices on an arbitrary vector (x, y) . The basis was therefore a pair of unit vectors pointing in the x and y directions. In most of the examples we will be considering in this course, we will use sets of atomic orbitals as basis functions for matrix representations. Don't worry too much if these ideas seem a little abstract at the moment – they should become clearer in the next section when we look at some examples.

Before proceeding any further, we must check that a matrix representation of a group obeys all of the rules set out in the formal mathematical definition of a group.

1. The first rule is that the group must include the identity operation E (the 'do nothing' operation). We showed above that the matrix representative of the identity operation is simply the identity matrix. As a consequence, every matrix representation includes the appropriate identity matrix.
2. The second rule is that the combination of any pair of elements must also be an element of the group (the *group property*). If we multiply together any two matrix representatives, we should get a new matrix which is a representative of another symmetry operation of the group. In fact, matrix representatives multiply together to give new representatives in exactly the same way as symmetry operations combine according to the group multiplication table. For example, in the C_{3v} point group, we showed that the combined symmetry operation $C_3\sigma_v$ is equivalent to σ_v'' . In a matrix representation of the group, if the matrix representatives of C_3 and σ_v are multiplied together, the result will be the representative of σ_v'' .
3. The third rule states that every operation must have an inverse, which is also a member of the group. The combined effect of carrying out an operation and its inverse is the same as the identity operation. It is fairly easy to show that matrix representatives satisfy this criterion. For example, the inverse of a reflection is another reflection, identical to the first. In matrix terms we would therefore expect that a reflection matrix was its own inverse, and that two identical reflection matrices multiplied together would give the identity matrix. This turns out to be true, and can be verified using any of the reflection matrices in the examples above. The inverse of a rotation matrix is another rotation matrix corresponding to a rotation of the opposite sense to the first.
4. The final rule states that the rule of combination of symmetry elements in a group must be associative. This is automatically satisfied by the rules of matrix multiplication.

Example: a matrix representation of the C_{3v} point group (the ammonia molecule)

The first thing we need to do before we can construct a matrix representation is to choose a basis. For NH_3 , we will select a basis (s_N, s_1, s_2, s_3) that consists of the valence s orbitals on the nitrogen and the three hydrogen atoms. We need to consider what happens to this basis when it is acted on by each of the symmetry operations in the C_{3v} point group, and determine the matrices that would be required to produce the same effect. The basis set and the symmetry operations in the C_{3v} point group are summarized in the figure below.



The effects of the symmetry operations on our chosen basis are as follows:

$$\begin{aligned}
 E & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_1, s_2, s_3) \\
 C_3^+ & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_2, s_3, s_1) \\
 C_3^- & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_3, s_1, s_2) \\
 \sigma_v & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_1, s_3, s_2) \\
 \sigma'_v & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_2, s_1, s_3) \\
 \sigma''_v & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_3, s_2, s_1)
 \end{aligned}
 \tag{1.10.1}$$

By inspection, the matrices that carry out the same transformations are:


$$\begin{aligned}
 \Gamma(E) & (s_N, s_1, s_2, s_3) \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = (s_N, s_1, s_2, s_3) \\
 \Gamma(C_3^+) & (s_N, s_1, s_2, s_3) \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} = (s_N, s_2, s_3, s_1) \\
 \Gamma(C_3^-) & (s_N, s_1, s_2, s_3) \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix} = (s_N, s_3, s_1, s_2) \\
 \Gamma(\sigma_v) & (s_N, s_1, s_2, s_3) \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} = (s_N, s_1, s_3, s_2) \\
 \Gamma(\sigma'_v) & (s_N, s_1, s_2, s_3) \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = (s_N, s_2, s_1, s_3) \\
 \Gamma(\sigma''_v) & (s_N, s_1, s_2, s_3) \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} = (s_N, s_3, s_2, s_1)
 \end{aligned}
 \tag{1.10.2}$$

These six matrices therefore form a representation for the C_{3v} point group in the (s_N, s_1, s_2, s_3) basis. They multiply together according to the group multiplication table and satisfy all the requirements for a mathematical group.

We have written the vectors representing our basis as row vectors. This is important. If we had written them as column vectors, the corresponding transformation matrices would be the transposes of the matrices above, and would not reproduce the group multiplication table (try it as an exercise if you need to convince yourself).

Example: a matrix representation of the C_{2v} point group (the allyl radical)

In this example, we'll take as our basis a p orbital on each carbon atom (p_1, p_2, p_3) .



Note that the p orbitals are *perpendicular* to the plane of the carbon atoms (this may seem obvious, but if you're visualizing the basis incorrectly it will shortly cause you a not inconsiderable amount of confusion). The symmetry operations in the C_{2v} point group, and their effect on the three p orbitals, are as follows:

$$\begin{aligned}
 E & (p_1, p_2, p_3) \rightarrow (p_1, p_2, p_3) \\
 C_2 & (p_1, p_2, p_3) \rightarrow (-p_3, -p_2, -p_1) \\
 \sigma_v & (p_1, p_2, p_3) \rightarrow (-p_1, -p_2, -p_3) \\
 \sigma'_v & (p_1, p_2, p_3) \rightarrow (p_3, p_2, p_1)
 \end{aligned}
 \tag{1.10.3}$$

The matrices that carry out the transformation are

$$\begin{aligned}
 \Gamma(E) & (p_1, p_2, p_3) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = (p_1, p_2, p_3) \\
 \Gamma(C_2) & (p_1, p_2, p_3) \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix} = (-p_3, -p_2, -p_1) \\
 \Gamma(\sigma_v) & (p_1, p_2, p_3) \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} = (-p_1, -p_2, -p_3) \\
 \Gamma(\sigma'_v) & (p_1, p_2, p_3) \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} = (p_3, p_2, p_1)
 \end{aligned}
 \tag{1.10.4}$$

This page titled [1.10: Matrix Representations of Groups](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.11: Properties of Matrix Representations

Now that we've learnt how to create a matrix representation of a point group within a given basis, we will move on to look at some of the properties that make these representations so powerful in the treatment of molecular symmetry.

Similarity Transforms

Suppose we have a basis set $(x_1, x_2, x_3, \dots, x_n)$, and we have determined the matrix representatives for the basis in a given point group. There is nothing particularly special about the basis set we have chosen, and we could equally well have used any set of linear combinations of the original functions (provided the combinations were linearly independent). The matrix representatives for the two basis sets will certainly be different, but we would expect them to be related to each other in some way. As we shall show shortly, they are in fact related by a *similarity transform*. It will be far from obvious at this point why we would want to carry out such a transformation, but similarity transforms will become important later on when we use group theory to choose an optimal basis set with which to generate molecular orbitals.

Consider a basis set $(x'_1, x'_2, x'_3, \dots, x'_n)$, in which each basis function x'_i is a linear combination of our original basis $(x_1, x_2, x_3, \dots, x_n)$.

$$x'_j = \sum_i x_i c_{ji} = x_1 c_{j1} + x_2 c_{j2} + \dots \quad (11.1)$$

The c_{ji} appearing in the sum are coefficients; c_{ji} is the coefficient multiplying the original basis function x_i in the new linear combination basis function x'_j . We could also represent this transformation in terms of a matrix equation $\mathbf{x}' = \mathbf{x}C$:

$$(x'_1, x'_2, \dots, x'_n) = (x_1, x_2, \dots, x_n) \begin{pmatrix} c_{11} & c_{12} & \dots & c_{1n} \\ c_{21} & c_{22} & \dots & c_{2n} \\ \dots & \dots & \dots & \dots \\ c_{n1} & c_{n2} & \dots & c_{nn} \end{pmatrix} \quad (11.2)$$

Now we look at what happens when we apply a symmetry operation g to our two basis sets. If $\Gamma(g)$ and $\Gamma'(g)$ are matrix representatives of the symmetry operation in the \mathbf{x} and \mathbf{x}' bases, then we have:

$$\begin{aligned} g\mathbf{x}' &= \mathbf{x}'\Gamma'(g) \\ g\mathbf{x}C &= \mathbf{x}C\Gamma'(g) && \text{since } \mathbf{x}' = \mathbf{x}C \\ g\mathbf{x} &= \mathbf{x}C\Gamma'(g)C^{-1} && \text{multiplying on the right by } C^{-1} \text{ and using } CC^{-1} = I \\ &= \mathbf{x}\Gamma(g) \end{aligned} \quad (11.3)$$

We can therefore identify the similarity transform relating $\Gamma(g)$, the matrix representative in our original basis, to $\Gamma'(g)$, the representative in the transformed basis. The transform depends only on the matrix of coefficients used to transform the basis functions.

$$\Gamma(g) = C\Gamma'(g)C^{-1} \quad (11.4)$$

Also,

$$\Gamma'(g) = C^{-1}\Gamma(g)C \quad (11.5)$$

Characters of Representations

The trace of a matrix representative $\Gamma(g)$ is usually referred to as the *character* of the representation under the symmetry operation g . We will soon come to see that the characters of a matrix representation are often more useful than the matrix representatives themselves. Characters have several important properties.

1. The character of a symmetry operation is invariant under a similarity transform
2. Symmetry operations belonging to the same class have the same character in a given representation. Note that the character for a given class may be different in different representations, and that more than one class may have the same character.

Proofs of the above two statements are given in the Appendix.

This page titled [1.11: Properties of Matrix Representations](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon

request.

1.12: Reduction of Representations I

Let us now go back and look at the C_{3v} representation we derived in 10.1 in more detail. If we look at the matrices carefully we see that they all take the same block diagonal form (a square matrix is said to be **block diagonal** if all the elements are zero except for a set of submatrices lying along the diagonal).

$$\begin{array}{cccccc}
 \Gamma(E) & \Gamma(C_3^-) & \Gamma(C_3^+) & \Gamma(\sigma_v) & \Gamma(\sigma'_v) & \Gamma(\sigma''_v) \\
 \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \\
 \chi(E) = 4 & \chi(C_3^+) = 1 & \chi(C_3^-) = 1 & \chi(\sigma_v) = 2 & \chi(\sigma'_v) = 2 & \chi(\sigma''_v) = 2
 \end{array}$$

A block diagonal matrix can be written as the *direct sum* of the matrices that lie along the diagonal. In the case of the C_{3v} matrix representation, each of the matrix representatives may be written as the direct sum of a 1×1 matrix and a 3×3 matrix.

$$\Gamma^{(4)}(g) = \Gamma^{(1)}(g) \otimes \Gamma^{(3)}(g) \quad (1.12.1)$$

in which the bracketed superscripts denote the dimensionality of the matrices. Note that a direct sum is very different from ordinary matrix addition since it produces a matrix of higher dimensionality. A direct sum of two matrices of orders n and m is performed by placing the matrices to be summed along the diagonal of a matrix of order $n + m$ and filling in the remaining elements with zeroes.

The reason why this result is useful in group theory is that the two sets of matrices $\Gamma^{(1)}(g)$ and $\Gamma^{(3)}(g)$ also satisfy all of the requirements for a matrix representation. Each set contains the identity and an inverse for each member, and the members multiply together associatively according to the group multiplication table³. Recall that the basis for the original four-dimensional representation had the s orbitals (s_N, s_1, s_2, s_3) of ammonia as its basis. The first set of reduced matrices, $\Gamma^{(1)}(g)$, forms a one-dimensional representation with (s_N) as its basis. The second set, $\Gamma^{(3)}(g)$ forms a three-dimensional representation with the basis (s_1, s_2, s_3). Separation of the original representation into representations of lower dimensionality is called *reduction* of the representation. The two *reduced representations* are shown below.

g	E	C_3^+	C_3^-	σ_v	σ'_v	σ''_v	
$\Gamma^{(1)}(g)$	(1)	(1)	(1)	(1)	(1)	(1)	1D representation spanned by (s_N)
$\Gamma^{(3)}(g)$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$	3D representation spanned by (s_1, s_2, s_3)

The logical next step is to investigate whether or not the three dimensional representation $\Gamma^{(3)}(g)$ can be reduced any further. As it stands, the matrices making up this representation are not in block diagonal form (some of you may have noted that the matrices representing E and σ_v are block diagonal, but in order for a representation to be reducible *all* of the matrix representatives must be in the *same* block diagonal form) so the representation is not reducible. However, we can carry out a similarity transformation (see 10.1) to a new representation spanned by a new set of basis functions (made up of linear combinations of (s_1, s_2, s_3)), which *is* reducible. In this case, the appropriate (normalized) linear combinations to use as our new basis functions are

$$\begin{aligned}
 s'_1 &= \frac{1}{\sqrt{3}}(s_1 + s_2 + s_3) \\
 s'_2 &= \frac{1}{\sqrt{6}}(2s_1 - s_2 - s_3) \\
 s'_3 &= \frac{1}{\sqrt{2}}(s_2 - s_3)
 \end{aligned} \quad (1.12.2)$$



or in matrix form

$$\begin{aligned}
 (s'_1, s'_2, s'_3) &= (s_1, s_2, s_3) \begin{pmatrix} \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{6}} & 0 \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{2}} \end{pmatrix} \\
 \mathbf{x}' &= \mathbf{x} C
 \end{aligned} \tag{1.12.3}$$

The matrices in the new representation are found from $\Gamma'(g) = C^{-1}\Gamma(g)C$ to be

$$\Gamma^{(3)}(g) \begin{matrix} E & C_3^+ & C_3^- & \sigma_v & \sigma'_v & \sigma''_v \\ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix} \end{matrix} \tag{1.12.4}$$

We see that each matrix is now in block diagonal form, and the representation may be reduced into the direct sum of a 1 x 1 representation spanned by (s'_1) and a 2x2 representation spanned by (s'_2, s'_3) . The complete set of reduced representations obtained from the original 4D representation is:

E	C_3^+	C_3^-	σ_v	σ'_v	σ''_v	
(1)	(1)	(1)	(1)	(1)	(1)	1D representation spanned by (s_N)
(1)	(1)	(1)	(1)	(1)	(1)	1D representation spanned by (s'_1)
$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$	$\begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$	2D representation spanned by (s'_2, s'_3)

(1.12.5)

This is as far as we can go in reducing this representation. None of the three representations above can be reduced any further, and they are therefore called *irreducible representations*, of the point group. Formally, a representation is an irreducible representation if there is no similarity transform that can simultaneously convert all of the representatives into block diagonal form. The linear combination of basis functions that converts a matrix representation into block diagonal form, allowing reduction of the representation, is called a *symmetry adapted linear combination (SALC)*.

³ The 1x1 representation in which all of the elements are equal to 1 is sometimes called the *unfaithful representation*, since it satisfies the group properties in a fairly trivial way without telling us much about the symmetry properties of the group.

1.13: Irreducible representations and symmetry species

The two one-dimensional irreducible representations spanned by s_N and s'_1 are seen to be identical. This means that s_N and s'_1 have the 'same symmetry', transforming in the same way under all of the symmetry operations of the point group and forming bases for the same matrix representation. As such, they are said to belong to the same *symmetry species*. There are a limited number of ways in which an arbitrary function can transform under the symmetry operations of a group, giving rise to a limited number of symmetry species. Any function that forms a basis for a matrix representation of a group must transform as one of the symmetry species of the group. The irreducible representations of a point group are labeled according to their symmetry species as follows:

- i. 1D representations are labeled A or B , depending on whether they are symmetric (character +1) or antisymmetric (character -1) under rotation about the principal axis.
- ii. 2D representations are labeled E , 3D representations are labeled T .
- iii. In groups containing a center of inversion, g and u labels (from the German *gerade* and *ungerade*, meaning symmetric and antisymmetric) denote the character of the irreducible representation under inversion (+1 for g , -1 for u)
- iv. In groups with a horizontal mirror plane but no center of inversion, the irreducible representations are given prime and double prime labels to denote whether they are symmetric (character +1) or antisymmetric (character -1) under reflection in the plane.
- v. If further distinction between irreducible representations is required, subscripts 1 and 2 are used to denote the character with respect to a C_2 rotation perpendicular to the principal axis, or with respect to a vertical reflection if there are no C_2 rotations.

The 1D irreducible representation in the C_{3v} point group is symmetric (has character +1) under all the symmetry operations of the group. It therefore belongs to the irreducible representation A_1 . The 2D irreducible representation has character 2 under the identity operation, -1 under rotation, and 0 under reflection, and belongs to the irreducible representation E .

Sometimes there is confusion over the relationship between a function f and its irreducible representation, but it is quite important that you understand the connection. There are several different ways of stating the relationship. For example, the following statements all mean the same thing:

- " f has A_2 symmetry"
- " f transforms as A_2 "
- " f has the same symmetry"

This page titled [1.13: Irreducible representations and symmetry species](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.14: Character Tables

A character table summarizes the behavior of all of the possible irreducible representations of a group under each of the symmetry operations of the group. The character table for C_{3v} is shown below.

$C_{3v}, 3m$	E	$2C_3$	$3\sigma_v$	$h = 6$
A_1	1	1	1	$z, z^2, x^2 + y^2$
A_2	1	1	-1	R_z
E	2	-1	0	$(x, y), (xy, x^2 + y^2), (xz, yz), (R_x, R_y)$

(1.14.1)

The various sections of the table are as follows:

- i. The first element in the table gives the name of the point group, usually in both Schoenflies (C_{3v}) and Hermann-Mauguin ($3m$) notation.
- ii. Along the first row are the symmetry operations of the group, E , $2C_3$ and $3\sigma_v$, followed by the order of the group. Because operations in the same class have the same character, symmetry operations are grouped into classes in the character table and not listed separately.
- iii. In the first column are the irreducible representations of the group. In C_{3v} the irreducible representations are A_1 , A_2 and E (the representation we considered above spans $2A_1 + E$).
- iv. The characters of the irreducible representations under each symmetry operation are given in the bulk of the table.
- v. The final column of the table lists a number of functions that transform as the various irreducible representations of the group. These are the Cartesian axes (x, y, z) , the Cartesian products $(z^2, x^2 + y^2, xy, yz)$, and the rotations (R_x, R_y, R_z) .

The functions listed in the final column of the table are important in many chemical applications of group theory, particularly in spectroscopy. For example, by looking at the transformation properties of x , y and z (sometimes given in character tables as T_x , T_y , T_z) we can discover the symmetry of translations along the x , y , and z axes. Similarly, R_x , R_y and R_z represent rotations about the three Cartesian axes. As we shall see later, the transformation properties of x , y , and z can also be used to determine whether or not a molecule can absorb a photon of x -, y -, or z -polarized light and undergo a spectroscopic transition. The Cartesian products play a similar role in determining selection rules for Raman transitions, which involve two photons.

Character tables for common point groups are given in Appendix B.

A simple way to determine the characters of a representation

In many applications of group theory, we only need to know the characters of the representative matrices, rather than the matrices themselves. Luckily, when each basis function transforms as a 1D irreducible representation (which is true in many cases of interest) there is a simple shortcut to determining the characters without having to construct the entire matrix representation. All we have to do is to look at the way the individual basis functions transform under each symmetry operation. For a given operation, step through the basis functions as follows:

- i. Add 1 to the character if the basis function is unchanged by the symmetry operation (i.e. the basis function is mapped onto itself);
- ii. Add -1 to the character if the basis function changes sign under the symmetry operation (i.e the basis function is mapped onto minus itself);
- iii. Add 0 to the character if the basis function moves when the symmetry operation is applied (i.e the basis function is mapped onto something different from itself).

Try this for the s orbital basis we have been using for the C_{3v} group. You should find you get the same characters as we obtained from the traces of the matrix representatives.

We can also work out the characters fairly easily when two basis functions transform together as a 2D irreducible representation. For example, in the C_{3v} point group x and y axes transform together as E . If we carry out a rotation about z by an angle θ , our x and y axes are transformed onto new axes x' and y' . However, the new axes can each be written as a linear combination of our original x and y axes. Using the rotation matrices introduced in Section 9, we see that:

$$\begin{aligned} x' &= \cos \theta x + \sin \theta y \\ y' &= -\sin \theta x + \cos \theta y \end{aligned} \quad (1.14.2)$$

For one-dimensional irreducible representations we asked if a basis function/axis was mapped onto itself, minus itself, or something different. For two-dimensional irreducible representations we need to ask how much of the 'old' axis is contained in the new one. From the above we see that the x' axis contains a contribution $\cos\theta$ from the x axis, and the y' axis contains a contribution $\cos\theta$ from the y axis. The characters of the x and y axes under a rotation through θ are therefore $\cos\theta$, and the overall character of the E irreducible representation is therefore $\cos\theta + \cos\theta = 2 \cos\theta$. For a C_3 rotation through 120 degrees, the character of the E irreducible representation is therefore $2 \cos 120^\circ = -1$.

In general, when an axis is rotated by an angle θ by a symmetry operation, its contribution to the character for that operation is $\cos\theta$.

Irreducible representations with complex characters

In many cases (see Appendix B), the characters for rotations C_n and improper rotations S_n are complex numbers, usually expressed in terms of the quantity $\epsilon = \exp(2\pi i/n)$. It is fairly straightforward to reconcile this with the fact that in chemistry we are generally using group theory to investigate physical problems in which all quantities are real. It turns out that whenever our basis spans an irreducible representation whose characters are complex, it will also span a second irreducible representation whose characters are the complex conjugates of the first irreducible representation i.e. complex irreducible representations occur in pairs. According to the strict mathematics of group theory, each irreducible representation in the pair should be considered as a separate representation. However, when applying such irreducible representations in physical problems, we add the characters for the two irreducible representations together to get a single irreducible representation whose characters are real.

As an example, the 'correct' character table for the group C_3 takes the form:

$$\begin{array}{c|ccc}
 C_3 & E & C_3 & C_3^2 \\
 \hline
 A & 1 & 1 & 1 \\
 \hline
 E & \left\{ \begin{array}{ccc} 1 & \epsilon & \epsilon^* \\ 1 & \epsilon^* & \epsilon \end{array} \right\} & &
 \end{array} \tag{1.14.3}$$

Where $\epsilon = \exp(2\pi i/3)$. However, as chemists we would usually combine the two parts of the E irreducible representation to give:

$$\begin{array}{c|ccc}
 C_3 & E & C_3 & C_3^2 \\
 \hline
 A & 1 & 1 & 1 \\
 \hline
 E & 2 & -1 & 1
 \end{array} \tag{1.14.4}$$

This page titled [1.14: Character Tables](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.15: Reduction of representations II

By making maximum use of molecular symmetry, we often greatly simplify problems involving molecular properties. For example, the formation of chemical bonds is strongly dependent on the atomic orbitals involved having the correct symmetries. To make full use of group theory in the applications we will be considering, we need to develop a little more ‘machinery’. Specifically, given a basis set (of atomic orbitals, for example) we need to find out:

1. How to determine the irreducible representations spanned by the basis functions
2. How to construct linear combinations of the original basis functions that transform as a given irreducible representation/symmetry species.

It turns out that both of these problems can be solved using something called the ‘**Great Orthogonality Theorem**’ (GOT for short). The GOT summarizes a number of orthogonality relationships implicit in matrix representations of symmetry groups, and may be derived in a somewhat qualitative fashion by considering these relationships in turn.

Some of you might find the next section a little hard going. In it, we will derive two important expressions that we can use to achieve the two goals we have set out above. It is not important that you understand every step in these derivations; they have mainly been included just so you can see where the equations come from. However, you *will* need to understand how to use the results. Hopefully you will not find this too difficult once we’ve worked through a few examples.

General concepts of Orthogonality


You are probably already familiar with the *geometric* concept of orthogonality. Two vectors are orthogonal if their dot product (i.e. the projection of one vector onto the other) is zero. An example of a pair of orthogonal vectors is provided by the **x** and **y** Cartesian unit vectors.

$$\mathbf{x}, \mathbf{y} = 0 \quad (1.15.1)$$



A consequence of the orthogonality of **x** and **y** is that any general vector in the *xy* plane may be written as a linear combination of these two *basis vectors*.

$$\mathbf{r} = a\mathbf{x} + b\mathbf{y} \quad (1.15.2)$$




Mathematical functions may also be orthogonal. Two functions, $f_1(x)$ and $f_2(x)$, are defined to be orthogonal if the integral over their product is equal to zero i.e.

$$\int f_1(x)f_2(x)dx = \delta_{12} \quad (1.15.3)$$

This simply means that there must be ‘no overlap’ between orthogonal functions, which is the same as the orthogonality requirement for vectors, above. In the same way as for vectors, any general function may be written as a linear combination of a suitably chosen set of orthogonal *basis functions*. For example, the Legendre polynomials $P_n(x)$ form an orthogonal basis set for functions of one variable x .

$$f(x) = \sum_n c_n P_n(x) \quad (1.15.4)$$



Orthogonality relationships in Group Theory

The irreducible representations of a point group satisfy a number of orthogonality relationships:

1. If corresponding matrix elements in all of the matrix representatives of an irreducible representation are squared and added together, the result is equal to the order of the group divided by the dimensionality of the irreducible representation. i.e.

$$\sum_g \Gamma_k(g)_{ij} \Gamma_k(g)_{ij} = \frac{h}{d_k} \quad (1.15.5)$$

where k labels the irreducible representation, i and j label the row and column position within the irreducible representation, h is the order of the group, and d_k is the order of the irreducible representation. e.g. The order of the group C_{3v} is 6. If we apply the above operation to the first element in the 2x2 (E) irreducible representation derived in Section 12, the result should be equal to $\frac{h}{d_k} = \frac{6}{2} = 3$. Carrying out this operation gives:

$$(1)^2 + \left(-\frac{1}{2}\right)^2 + \left(-\frac{1}{2}\right)^2 + (1)^2 + \left(-\frac{1}{2}\right)^2 + \left(-\frac{1}{2}\right)^2 = 1 + \frac{1}{4} + \frac{1}{4} + 1 + \frac{1}{4} + \frac{1}{4} = 3 \quad (1.15.6)$$

2. If instead of summing the squares of matrix elements in an irreducible representation, we sum the product of two different elements from within each matrix, the result is equal to zero. i.e.

$$\sum_g \Gamma_k(g)_{ij} \Gamma_k(g)_{i'j'} = 0 \quad (1.15.7)$$

where $i \neq i'$ and/or $j \neq j'$. E.g. if we perform this operation using the two elements in the first row of the 2D irreducible representation used in 1, we get:

$$(1)(0) + \left(-\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}\right) + \left(-\frac{1}{2}\right)\left(-\frac{\sqrt{3}}{2}\right) + (1)(0) + \left(-\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}\right) + \left(-\frac{1}{2}\right)\left(-\frac{\sqrt{3}}{2}\right) = 0 + \frac{\sqrt{3}}{4} - \frac{\sqrt{3}}{4} + 0 - \frac{\sqrt{3}}{4} + \frac{\sqrt{3}}{4} = 0 \quad (1.15.8)$$

3. If we sum the product of two elements from the matrices of two *different* irreducible representations k and m , the result is equal to zero. i.e.

$$\sum_g \Gamma_k(g)_{ij} \Gamma_m(g)_{i'j'} = 0 \quad (1.15.9)$$

where there is now no restriction on the values of the indices i, i', j, j' (apart from the rather obvious restriction that they must be less than or equal to the dimensions of the irreducible representation). e.g. Performing this operation on the first elements of the A_1 and E irreducible representations we derived for C_{3v} gives:

$$(1)(1) + (1)\left(-\frac{1}{2}\right) + (1)\left(-\frac{1}{2}\right) + (1)(1) + (1)\left(-\frac{1}{2}\right) + (1)\left(-\frac{1}{2}\right) = 1 - \frac{1}{2} - \frac{1}{2} + 1 - \frac{1}{2} - \frac{1}{2} = 0 \quad (1.15.10)$$

We can combine these three results into one general equation, the Great Orthogonality Theorem⁴.

$$\sum_g \Gamma_k(g)_{ij} \Gamma_m(g)_{i'j'} = \frac{h}{\sqrt{d_k d_m}} \delta_{km} \delta_{ii'} \delta_{jj'} \quad (1.15.11)$$

For most applications we do not actually need the full Great Orthogonality Theorem. A little mathematical trickery transforms Equation 1.15.11 into the 'Little Orthogonality Theorem' (or LOT), which is expressed in terms of the characters of the irreducible representations rather than the irreducible representations themselves.

$$\sum_g \chi_k(g) \chi_m(g) = h \delta_{km} \quad (1.15.12)$$

Since the characters for two symmetry operations in the same class are the same, we can also rewrite the sum over symmetry operations as a sum over classes.

$$\sum_C n_C \chi_k(C) \chi_m(C) = h \delta_{km} \quad (1.15.13)$$

where n_C is the number of symmetry operations in class C .

In all of the examples we've considered so far, the characters have been real. However, this is not necessarily true for all point groups, so to make the above equations completely general we need to include the possibility of imaginary characters. In this case we have:

$$\sum_C n_C \chi_k^*(C) \chi_m(C) = h \delta_{km} \quad (1.15.14)$$

where $\chi_k^*(C)$ is the complex conjugate of $\chi_k(C)$. Equation 1.15.14 is of course identical to Equation 1.15.13 when all the characters are real.

Using the LOT to Determine the Irreducible Representations Spanned by a Basis

In Section 12 we discovered that we can often carry out a similarity transform on a general matrix representation so that all the representatives end up in the same block diagonal form. When this is possible, each set of submatrices also forms a valid matrix representation of the group. If none of the submatrices can be reduced further by carrying out another similarity transform, they are said to form an irreducible representation of the point group. An important property of matrix representatives is that their character is invariant under a similarity transform. This means

that the character of the original representatives must be equal to the sum of the characters of the irreducible representations into which the representation is reduced. e.g. if we consider the representative for the C_3^- symmetry operation in our NH_3 example, we have:

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \xrightarrow{\text{similarity transform}} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 0 & 0 & \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} = (1) \otimes (1) \otimes \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} \quad (1.15.15)$$

$\chi = 1$ $\chi = 1$ $\chi = 1 + 1 - 1 = 1$

It follows that we can write the characters for a general representation $\Gamma(g)$ in terms of the characters of the irreducible representations $\Gamma_k(g)$ into which it can be reduced.

$$\chi(g) = \sum_k a_k \chi_k(g) \quad (1.15.16)$$

where the coefficients a_k in the sum are the number of times each irreducible representation appears in the representation. This means that in order to determine the irreducible representations spanned by a given basis, all we have to do is determine the coefficients a_k in the above equation. This is where the **Little Orthogonality Theorem** comes in handy. If we take the LOT in the form of Equation 1.15.16 and multiply each side through by a_k , we get

$$\sum_g a_k \chi_k(g) \chi_m(g) = h a_k \delta_{km} \quad (1.15.17)$$

Summing both sides of the above equation over k gives

$$\sum_g \sum_k a_k \chi_k(g) \chi_m(g) = h \sum_k a_k \delta_{km} \quad (1.15.18)$$

We can use Equation 1.15.16 to simplify the left hand side of this equation. Also, the sum on the right hand side reduces to a_m because δ_{km} is only non-zero (and equal to 1) when $k = m$

$$\sum_g \chi(g) \chi_m(g) = h a_m \quad (1.15.19)$$

Dividing both sides through by h (the order of the group), gives us an expression for the coefficients a_m in terms of the characters $\chi(g)$ of the original representation and the characters $\chi_m(g)$ of the m^{th} irreducible representation.

$$a_m = \frac{1}{h} \sum_g \chi(g) \chi_m(g) \quad (1.15.20)$$

We can of course write this as a sum over classes rather than a sum over symmetry operations.

$$a_m = \frac{1}{h} \sum_C n_C \chi(g) \chi_m(g) \quad (1.15.21)$$

As an example, in Section 12 we showed that the matrix representatives we derived for the C_{3v} group could be reduced into two irreducible representations of A_1 symmetry and one of E symmetry. i.e. $\Gamma = 2A_1 + E$. We could have obtained the same result using Equation 1.15.21. The characters for our original representation and for the irreducible representations of the C_{3v} point group (A_1 , A_2 and E) are given in the table below.

C_{3v}	E	$2C_3$	$3\sigma_v$
χ	4	1	2
$\chi(A_1)$	1	1	1
$\chi(A_2)$	1	1	-1
$\chi(E)$	2	-1	0

(1.15.22)

From Equation 1.15.21, the number of times each irreducible representation occurs for our chosen basis (s_n, s_1, s_2, s_3) is therefore

$$\begin{aligned} a(A_1) &= \frac{1}{6} (1x4x1 + 2x1x1 + 3x2x1) = 2 \\ a(A_2) &= \frac{1}{6} (1x4x1 + 2x1x1 + 3x2x-1) = 0 \\ a(E) &= \frac{1}{6} (1x4x2 + 2x1x-1 + 3x2x0) = 1 \end{aligned} \quad (1.15.23)$$

i.e. Our basis is spanned by $2A_1 + E$, as we found before.

⁴The δ_{ij} appearing in Equation 1.15.11 are called Dirac delta functions. They are equal to 1 if $i = j$ and 0 otherwise.

This page titled 1.15: Reduction of representations II is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.16: Symmetry Adapted Linear Combinations (SALCs)

Once we know the irreducible representations spanned by an arbitrary basis set, we can work out the appropriate linear combinations of basis functions that transform the matrix representatives of our original representation into block diagonal form (i.e. the *symmetry adapted linear combinations*). Each of the SALCs transforms as one of the irreducible representations of the reduced representation. We have already seen this in our NH_3 example. The two linear combinations of A_1 symmetry were s_N and $s_1 + s_2 + s_3$, both of which are symmetric under all the symmetry operations of the point group. We also chose another pair of functions, $2s_1 - s_2 - s_3$ and $s_2 - s_3$, which together transform as the symmetry species E .

To find the appropriate SALCs to reduce a matrix representation, we use *projection operators*. You will be familiar with the idea of operators from quantum mechanics. The operators we will be using here are not quantum mechanical operators, but the basic principle is the same. The projection operator to generate a SALC that transforms as an irreducible representation k is $\sum_g \chi_k(g)g$. Each term in the sum means ‘apply the symmetry operation g and then multiply by the character of g in irreducible representation k ’. Applying this operator to each of our original basis functions in turn will generate a complete set of SALCs, i.e. to transform a basis function f_i into a SALC f'_i , we use

$$f'_i = \sum_g \chi_k(g)gf_i \quad (16.1)$$

The way in which this operation is carried out will become much more clear if we work through an example. We can break down the above equation into a fairly straightforward ‘recipe’ for generating SALCs:

1. Make a table with columns labeled by the basis functions and rows labeled by the symmetry operations of the molecular point group. In the columns, show the effect of the symmetry operations on the basis functions (this is the gf_i part of Equation 16.1).
2. For each irreducible representation in turn, multiply each member of the table by the character of the appropriate symmetry operation (we now have $\chi_k(g)gf_i$ for each operation). Summing over the columns (symmetry operations) generates all the SALCs that transform as the chosen irreducible representation.
3. Normalize the SALCs.

Earlier (see Section 10), we worked out the effect of all the symmetry operations in the C_{3v} point group on the (s_N, s_1, s_2, s_3) basis.

$$\begin{array}{ll}
 E & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_1, s_2, s_3) \\
 C_3^+ & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_2, s_3, s_1) \\
 C_3^- & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_3, s_1, s_2) \\
 \sigma_v & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_1, s_3, s_2) \\
 \sigma'_v & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_2, s_1, s_3) \\
 \sigma''_v & (s_N, s_1, s_2, s_3) \rightarrow (s_N, s_3, s_2, s_1)
 \end{array} \quad (16.2)$$

This is all we need to construct the table described in 1. above.

	s_N	s_1	s_2	s_3	
E	s_N	s_1	s_2	s_3	
C_3^+	s_N	s_2	s_3	s_1	
C_3^-	s_N	s_3	s_1	s_2	
σ_v	s_N	s_1	s_3	s_2	
σ'_v	s_N	s_2	s_1	s_3	
σ''_v	s_N	s_3	s_2	s_1	

(16.3)

To determine the SALCs of A_1 symmetry, we multiply the table through by the characters of the A_1 irreducible representation (all of which take the value 1). Summing the columns gives

$$\begin{aligned}
 s_N + s_N + s_N + s_N + s_N + s_N &= 6s_N \\
 s_1 + s_2 + s_3 + s_1 + s_2 + s_3 &= 2(s_1 + s_2 + s_3) \\
 s_2 + s_3 + s_1 + s_3 + s_1 + s_2 &= 2(s_1 + s_2 + s_3) \\
 s_3 + s_1 + s_2 + s_2 + s_1 + s_3 &= 2(s_1 + s_2 + s_3)
 \end{aligned}
 \tag{16.4}$$

Apart from a constant factor (which doesn't affect the functional form and therefore doesn't affect the symmetry properties), these are the same as the combinations we determined earlier. Normalizing gives us two SALCs of A_1 symmetry.

$$\begin{aligned}
 \phi_1 &= s_N \\
 \phi_2 &= \frac{1}{\sqrt{3}}(s_1 + s_2 + s_3)
 \end{aligned}
 \tag{16.5}$$

We now move on to determine the SALCs of E symmetry. Multiplying the table above by the appropriate characters for the E irreducible representation gives

	s_N	s_1	s_2	s_3
E	$2s_N$	$2s_1$	$2s_2$	$2s_3$
C_3^+	$-s_N$	$-s_2$	$-s_3$	$-s_1$
C_3^-	$-s_N$	$-s_3$	$-s_1$	$-s_2$
σ_v	0	0	0	0
σ'_v	0	0	0	0
σ''_v	0	0	0	0

(16.6)

Summing the columns yields

$$\begin{aligned}
 2s_N - s_N - s_N &= 0 \\
 2s_1 - s_2 - s_3 & \\
 2s_2 - s_3 - s_1 & \\
 2s_3 - s_1 - s_2 &
 \end{aligned}
 \tag{16.7}$$

We therefore get three SALCs from this procedure. This is a problem, since the number of SALCs must match the dimensionality of the irreducible representation, in this case two. Put another way, we should end up with four SALCs in total to match our original number of basis functions. Added to our two SALCs of A_1 symmetry, three SALCs of E symmetry would give us five in total.

The resolution to our problem lies in the fact that the three SALCs above are not linearly independent. Any one of them can be written as a linear combination of the other two e.g. $(2s_1 - s_2 - s_3) = -(2s_2 - s_3s_1) - (2s_3 - s_1 - s_2)$. To solve the problem, we can either throw away one of the SALCs, or better, make two linear combinations of the three SALCs that are orthogonal to each other.⁵ e.g. if we take $2s_1 - s_2 - s_3$ as one of our SALCs and find an orthogonal combination of the other two (which turns out to be their difference), we have (after normalization)

$$\begin{aligned}
 \phi_3 &= \frac{1}{\sqrt{6}}(2s_1 - s_2 - s_3) \\
 \phi_4 &= \frac{1}{\sqrt{2}}(s_2 - s_3)
 \end{aligned}
 \tag{16.8}$$

These are the same linear combinations used in Section 12.

We now have all the machinery we need to apply group theory to a range of chemical problems. In our first application, we will learn how to use molecular symmetry and group theory to help us understand chemical bonding.

⁵ If we write the coefficients of s_1 , s_2 and s_3 for each SALC as a vector (a_1, a_2, a_3) , then when two SALCs are orthogonal, the dot product of their coefficient vectors $(a_1, a_2, a_3) \cdot (b_1, b_2, b_3) = (a_1b_1 + a_2b_2 + a_3b_3)$ is equal to zero.

This page titled [1.16: Symmetry Adapted Linear Combinations \(SALCs\)](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.17: Determining whether an Integral can be Non-zero

As we continue with this course, we will discover that there are many times when we would like to know whether a particular integral is necessarily zero, or whether there is a chance that it may be non-zero. We can often use group theory to differentiate these two cases.

You will have already used symmetry properties of functions to determine whether or not a one-dimensional integral is zero. For example, $\sin(x)$ is an 'odd' function (antisymmetric with respect to reflection through the origin), and it follows from this that

$$\int_{-\infty}^{\infty} \cos(x) dx = 0 \quad (1.17.1)$$

In general, an integral between these limits for any other odd function will also be zero.

In the general case we may have an integral of more than one dimension. The key to determining whether a general integral is necessarily zero lies in the fact that because an integral is just a number, it must be invariant to any symmetry operation. For example, bonding in a diatomic (see next section) depends on the presence of a non-zero overlap between atomic orbitals on adjacent atoms, which may be quantified by an *overlap integral*. You would not expect the bonding in a molecule to change if you rotated the molecule through some angle θ , so the integral must be invariant to rotation, and indeed to any other symmetry operation.

In group theoretical terms, for an integral to be non-zero, *the integrand must transform as the totally symmetric irreducible representation in the appropriate point group*. In practice, the integrand may not transform as a single irreducible representation, but it **must include** the totally symmetric irreducible representation. These ideas should become more clear in the next section.

Note

It should be noted that even when the irreducible representations spanned by the integrand do include the totally symmetric irreducible representation, it is still possible for the integral to be zero. All group theory allows us to do is identify integrals that are *necessarily* zero based on the symmetry (or lack thereof) of the integrand.

This page titled [1.17: Determining whether an Integral can be Non-zero](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.


1.18: Bonding in Diatomics

You will already be familiar with the idea of constructing molecular orbitals from linear combinations of atomic orbitals from previous courses covering bonding in diatomic molecules. By considering the symmetries of s and p orbitals on two atoms, we can form **bonding** and **antibonding** combinations labeled as having either σ or π symmetry depending on whether they resemble s or p orbitals when viewed along the bond axis (see diagram below). In all of the cases shown, only atomic orbitals that have the same symmetry when viewed along the bond axis z can form a chemical bond e.g. two s orbitals, two p_z orbitals, or an s and a p_z can form a bond, but a p_z and a p_x or an s and a p_x or a p_y cannot. It turns out that the rule that determines whether or not two atomic orbitals can bond is that *they must belong to the same symmetry species within the point group of the molecule*.

We can prove this mathematically for two atomic orbitals ϕ_i and ϕ_j by looking at the overlap integral between the two orbitals.

$$S_{ij} = \langle \phi_i | \phi_j \rangle = \int \phi_i^* \phi_j d\tau \quad (18.1) \quad (1.18.1)$$

In order for bonding to be possible, this integral must be non-zero. The product of the two functions ϕ_1 and ϕ_2 transforms as the direct product of their symmetry species i.e. $\Gamma_{12} = \Gamma_1 \otimes \Gamma_2$. As explained above, for the overlap integral to be non-zero, Γ_{12} must contain the totally symmetric irreducible representation (A_{1g} for a homonuclear diatomic, which belongs to the point group $D_{\infty h}$). As it happens, this is only possible if ϕ_1 and ϕ_2 belong to the same irreducible representation. These ideas are summarized for a diatomic in the table below.



First Atomic Orbital	Second Atomic Orbital	$\Gamma_1 \otimes \Gamma_2$	Overlap Integral	Bonding?
$s (A_{1g})$	$s (A_{1g})$	A_{1g}	Non-zero	Yes
$s (A_{1g})$	$p_x (E_{1u})$	E_{1u}	Zero	No
$s (A_{1g})$	$p_z (A_{1u})$	A_{1u}	Zero	No
$p_x (E_{1u})$	$p_x (E_{1u})$	$A_{1g} + A_{2g} + E_{2g}$	Non-zero	Yes
$p_x (E_{1u})$	$p_z (A_{1u})$	E_{1g}	Zero	No
$p_z (A_{1u})$	$p_z (A_{1u})$	A_{1g}	Non-zero	Yes

(18.2)

This page titled [1.18: Bonding in Diatomics](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.19: Bonding in Polyatomics- Constructing Molecular Orbitals from SALCs

In the previous section we showed how to use symmetry to determine whether two atomic orbitals can form a chemical bond. How do we carry out the same procedure for a polyatomic molecule, in which many atomic orbitals may combine to form a bond? Any SALCs of the same symmetry could potentially form a bond, so all we need to do to construct a molecular orbital is take a linear combination of all the SALCs of the same symmetry species. The general procedure is:

1. Use a basis set consisting of valence atomic orbitals on each atom in the system.
2. Determine which irreducible representations are spanned by the basis set and construct the SALCs that transform as each irreducible representation.
3. Take linear combinations of irreducible representations of the same symmetry species to form the molecular orbitals. E.g. in our NH_3 example we could form a molecular orbital of A_1 symmetry from the two SALCs that transform as A_1 ,

$$\begin{aligned}\Psi(A_1) &= c_1\phi_1 + c_2\phi_2 \\ &= c_1s_N + c_2\frac{1}{\sqrt{3}}(s_1 + s_2 + s_3)\end{aligned}\tag{19.1}$$

Unfortunately, this is as far as group theory can take us. It can give us the functional form of the molecular orbitals but it cannot determine the coefficients c_1 and c_2 . To go further and obtain the expansion coefficients and orbital energies, we must turn to quantum mechanics. The material we are about to cover will be repeated in greater detail in later courses on quantum mechanics and valence, but they are included here to provide you with a complete reference on how to construct molecular orbitals and determine their energies.

This page titled [1.19: Bonding in Polyatomics- Constructing Molecular Orbitals from SALCs](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.20: Calculating Orbital Energies and Expansion Coefficients

Calculation of the orbital energies and expansion coefficients is based on the *variation principle*, which states that any approximate wavefunction must have a higher energy than the true wavefunction. This follows directly from the fairly common-sense idea that in general any system tries to minimize its energy. If an ‘approximate’ wavefunction had a lower energy than the ‘true’ wavefunction, we would expect the system to try and adopt this ‘approximate’ lower energy state, rather than the ‘true’ state. That all approximations to the true wavefunction must have a higher energy than the true wavefunction is the only scenario that makes physical sense. A mathematical proof of the variation principle is given in the Appendix.

We apply the variation principle as follows:

Molecular energy levels, or orbital energies, are eigenvalues of the molecular Hamiltonian \hat{H} . Using a standard result from quantum mechanics, it follows that the energy E of a molecular orbital Ψ is

$$E = \frac{\langle \Psi | \hat{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} \quad (\text{unnormalized } \Psi) \quad (1.20.1)$$

or $E = \langle \Psi | \hat{H} | \Psi \rangle$ (normalized Ψ , for which $\langle \Psi | \Psi \rangle = 1$)

If the true wavefunction has the lowest energy, then to find the closest approximation we can to the true wavefunction, all we have to do is find the coefficients in our expansion of SALCs that minimize the energy in the above expressions. In practice, we substitute our wavefunction and minimize the resulting expression with respect to the coefficients. To show how this is done, we’ll use our NH_3 wavefunction of A_1 symmetry from the previous section. Substituting into Equation 1.20.1 gives:

$$\begin{aligned} E &= \frac{\langle c_1 \phi_1 + c_2 \phi_2 | \hat{H} | c_1 \phi_1 + c_2 \phi_2 \rangle}{\langle c_1 \phi_1 + c_2 \phi_2 | c_1 \phi_1 + c_2 \phi_2 \rangle} \\ &= \frac{\langle c_1 \phi_1 | \hat{H} | c_1 \phi_1 \rangle + \langle c_1 \phi_1 | \hat{H} | c_2 \phi_2 \rangle + \langle c_2 \phi_2 | \hat{H} | c_1 \phi_1 \rangle + \langle c_2 \phi_2 | \hat{H} | c_2 \phi_2 \rangle}{\langle c_1 \phi_1 | c_1 \phi_1 \rangle + \langle c_1 \phi_1 | c_2 \phi_2 \rangle + \langle c_2 \phi_2 | c_1 \phi_1 \rangle + \langle c_2 \phi_2 | c_2 \phi_2 \rangle} \\ &= \frac{c_1^2 \langle \phi_1 | \hat{H} | \phi_1 \rangle + c_1 c_2 \langle \phi_1 | \hat{H} | \phi_2 \rangle + c_2 c_1 \langle \phi_2 | \hat{H} | \phi_1 \rangle + c_2^2 \langle \phi_2 | \hat{H} | \phi_2 \rangle}{c_1^2 \langle \phi_1 | \phi_1 \rangle + c_1 c_2 \langle \phi_1 | \phi_2 \rangle + c_2 c_1 \langle \phi_2 | \phi_1 \rangle + c_2^2 \langle \phi_2 | \phi_2 \rangle} \end{aligned} \quad (1.20.2)$$

If we now define a Hamiltonian matrix element $H_{ij} = \langle \phi_i | \hat{H} | \phi_j \rangle$ and an overlap integral $S_{ij} = \langle \phi_i | \phi_j \rangle$ and note that $H_{ij} = H_{ji}$ and $S_{ij} = S_{ji}$, this simplifies to

$$E = \frac{c_1^2 H_{11} + 2c_1 c_2 H_{12} + c_2^2 H_{22}}{c_1^2 S_{11} + 2c_1 c_2 S_{12} + c_2^2 S_{22}} \quad (1.20.3)$$

To get this into a simpler form for carrying out the energy minimization, we multiply both sides through by the denominator to give

$$E(c_1^2 S_{11} + 2c_1 c_2 S_{12} + c_2^2 S_{22}) = c_1^2 H_{11} + 2c_1 c_2 H_{12} + c_2^2 H_{22} \quad (1.20.4)$$

Now we need to minimize the energy with respect to c_1 and c_2 , i.e., we require

$$\frac{\partial E}{\partial c_1} = 0 \quad (1.20.5)$$

and

$$\frac{\partial E}{\partial c_2} = 0 \quad (1.20.6)$$

If we differentiate the above equation through separately by c_1 and c_2 and apply this condition, we will end up with two equations in the two unknowns c_1 and c_2 , which we can solve to determine the coefficients and the energy.

Differentiating Equation 1.20.4 with respect to c_1 (via the product rule of differentiation) gives

$$\frac{\partial E}{\partial c_1} (c_1^2 S_{11} + 2c_1 c_2 S_{12} + c_2^2 S_{22}) + E(2c_1 S_{11} + 2c_2 S_{12}) = 2c_1 H_{11} + 2c_2 H_{12} \quad (1.20.7)$$

Differentiating Equation 1.20.4 with respect to c_2 gives

$$\frac{\partial E}{\partial c_2}(c_1^2 S_{11} + 2c_1 c_2 S_{12} + c_2^2 S_{22}) + E(2c_1 S_{12} + 2c_2 S_{22}) = 2c_1 H_{12} + 2c_2 H_{22} \quad (1.20.8)$$

Because

$$\frac{\partial E}{\partial c_1} = \frac{\partial E}{\partial c_2} = 0 \quad (1.20.9)$$

the first term on the left hand side of both equations is zero, leaving us with

$$\begin{aligned} E(2c_1 S_{11} + 2c_2 S_{12}) &= 2c_1 H_{11} + 2c_2 H_{12} \\ E(2c_1 S_{12} + 2c_2 S_{22}) &= 2c_1 H_{12} + 2c_2 H_{22} \end{aligned} \quad (1.20.10)$$

These are normally rewritten slightly, in the form

$$\begin{aligned} c_1(H_{11} - ES_{11}) + c_2(H_{12} - ES_{12}) &= 0 \\ c_1(H_{12} - ES_{12}) + c_2(H_{22} - ES_{22}) &= 0 \end{aligned} \quad (1.20.11)$$

Equations 1.20.11 are known as the **secular equations** and are the set of equations we need to solve to determine c_1 , c_2 , and E . In the general case (derived in the Appendix), when our wavefunction is a linear combination of N SALCs (i.e. $\Psi = \sum_{i=1}^N c_i \phi_i$) we get N equations in N unknowns, with the k^{th} equation given by

$$\sum_{i=1}^N c_i (H_{ki} - ES_{ki}) = 0 \quad (1.20.12)$$

Note that we can use any basis functions we like together with the linear variation method described here to construct approximate molecular orbitals and determine their energies, but choosing to use SALCs simplifies things considerably when the number of basis functions is large. An arbitrary set of N basis functions leads to a set of N equations in N unknowns, which must be solved simultaneously. Converting the basis into a set of SALCs separates the equations into several smaller sets of secular equations, one for each irreducible representation, which can be solved independently. It is usually easier to solve several sets of secular equations of lower dimensionality than one set of higher dimensionality.

This page titled [1.20: Calculating Orbital Energies and Expansion Coefficients](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.21: Solving the Secular Equations

Matrix formulation of a set of linear equations

As we have seen already, any set of linear equations may be rewritten as a matrix equation $A\mathbf{x} = \mathbf{b}$. Linear equations are classified as *simultaneous linear equations* or *homogeneous linear equations*, depending on whether the vector \mathbf{b} on the RHS of the equation is non-zero or zero.

For a set of simultaneous linear equations (non-zero \mathbf{b}) it is fairly apparent that if a unique solution exists, it can be found by multiplying both sides by the inverse matrix A^{-1} (since $A^{-1}A$ on the left hand side is equal to the identity matrix, which has no effect on the vector \mathbf{x})

$$\begin{aligned} A\mathbf{x} &= \mathbf{b} \\ A^{-1}A\mathbf{x} &= A^{-1}\mathbf{b} \\ \mathbf{x} &= A^{-1}\mathbf{b} \end{aligned} \quad (1.21.1)$$

In practice, there are easier matrix methods for solving simultaneous equations than finding the inverse matrix, but these need not concern us here. In Section 8.4, we discovered that in order for a matrix to have an inverse, **it must have a non-zero determinant**. Since A^{-1} must exist in order for a set of simultaneous linear equations to have a solution, this means that the determinant of the matrix A must be non-zero for the equations to be solvable.

The reverse is true for homogeneous linear equations. In this case the set of equations only has a solution if the determinant of A is equal to zero. The secular equations we want to solve are homogeneous equations, and we will use this property of the determinant to determine the molecular orbital energies. An important property of homogeneous equations is that if a vector \mathbf{x} is a solution, so is any multiple of \mathbf{x} , meaning that the solutions (the molecular orbitals) can be normalized without causing any problems.

Solving for the orbital energies and expansion coefficients

Recall the secular equations for the A_1 orbitals of NH_3 derived in the previous section

$$\begin{aligned} c_1(H_{11} - ES_{11}) + c_2(H_{12} - ES_{12}) &= 0 \\ c_1(H_{12} - ES_{12}) + c_2(H_{22} - ES_{22}) &= 0 \end{aligned} \quad (1.21.2)$$

where c_1 and c_2 are the coefficients in the linear combination of the SALCs $\phi_1 = s_N$ and $\phi_2 = \frac{1}{\sqrt{3}}(s_1 + s_2 + s_3)$ used to construct the molecular orbital. Writing this set of homogeneous linear equations in matrix form gives

$$\begin{pmatrix} H_{11} - ES_{11} & H_{12} - ES_{12} \\ H_{12} - ES_{12} & H_{22} - ES_{22} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (1.21.3)$$

For the equations to have a solution, the determinant of the matrix must be equal to zero. Writing out the determinant will give us a polynomial equation in E that we can solve to obtain the orbital energies in terms of the Hamiltonian matrix elements H_{ij} and overlap integrals S_{ij} . The number of energies obtained by ‘solving the secular determinant’ in this way is equal to the order of the matrix, in this case two.

The secular determinant for Equation 1.21.3 is (noting that $S_{11} = S_{22} = 1$ since the SALCs are normalized)

$$(H_{11} - E)(H_{22} - E) - (H_{12} - ES_{12})^2 = 0 \quad (1.21.4)$$

Expanding and collecting terms in E gives

$$E^2(1 - S_{12}^2) + E(2H_{12}S_{12} - H_{11} - H_{22}) + (H_{11}H_{22} - H_{12}^2) = 0 \quad (1.21.5)$$

which can be solved using the quadratic formula to give the energies of the two molecular orbitals.

$$E_{\pm} = \frac{-(2H_{12}S_{12} - H_{11} - H_{22}) \pm \sqrt{(2H_{12}S_{12} - H_{11} - H_{22})^2 - 4(1 - S_{12}^2)(H_{11}H_{22} - H_{12}^2)}}{2(1 - S_{12}^2)} \quad (1.21.6)$$

To obtain numerical values for the energies, we need to evaluate the integrals H_{11} , H_{22} , H_{12} , and S_{12} . This would be quite a challenge to do analytically, but luckily there are a number of computer programs that can be used to calculate the integrals. One

such program gives the following values.

$$\begin{aligned} H_{11} &= -26.0000 \text{ eV} \\ H_{22} &= -22.2216 \text{ eV} \\ H_{12} &= -29.7670 \text{ eV} \\ S_{12} &= 0.8167 \text{ eV} \end{aligned} \quad (1.21.7)$$

When we substitute these into our equation for the energy levels, we get:

$$\begin{aligned} E_+ &= 29.8336 \text{ eV} \\ E_- &= -31.0063 \text{ eV} \end{aligned} \quad (1.21.8)$$

We now have the orbital energies and the next step is to find the orbital coefficients. The coefficients for an orbital of energy E are found by substituting the energy into the secular equations and solving for the coefficients c_i . Since the two secular equations are not linearly independent (i.e. they are effectively only one equation), when we solve them to find the coefficients what we will end up with is the *relative* values of the coefficients. This is true in general: in a system with N coefficients, solving the secular equations will allow all N of the coefficients c_i to be obtained in terms of, say, c_1 . The absolute values of the coefficients are found by normalizing the wavefunction.

Since the secular equations for the orbitals of energy E_+ and E_- are not linearly independent, we can choose to solve either one of them to find the orbital coefficients. We will choose the first.

$$(H_{11} - E_{\pm})c_1 + (H_{12} - E_{\pm}S_{12})c_2 = 0 \quad (1.21.9)$$

For the orbital with energy $E_- = -31.0063$ eV, substituting numerical values into this equation gives

$$\begin{aligned} 5.0063c_1 - 4.4442c_2 &= 0 \\ c_2 &= 1.1265c_1 \end{aligned} \quad (1.21.10)$$

The molecular orbital is therefore

$$\Psi = c_1(\phi_1 + 1.1265\phi_2) \quad (1.21.11)$$

Normalizing to find the constant c_1 (by requiring $|\Psi| = 1$) gives

$$\begin{aligned} \Psi_1 &= 0.4933\phi_1 + 0.5557\phi_2 \\ &= 0.4933s_N + 0.3208(s_1 + s_2 + s_3) \quad (\text{substituting the SALCs for } \phi_1 \text{ and } \phi_2) \end{aligned} \quad (1.21.12)$$

For the second orbital, with energy $E_+ = 29.8336$ eV, the secular equation is

$$\begin{aligned} -55.8336c_1 - 54.1321c_2 &= 0 \\ c_2 &= -1.0314c_1 \end{aligned} \quad (1.21.13)$$

giving

$$\begin{aligned} \Psi_2 &= c_1(\phi_1 - 1.0314\phi_2) \\ &= 1.6242\phi_1 - 1.6752\phi_2 \quad (\text{after normalization}) \\ &= 1.6242s_N - 0.9672(s_1 + s_2 + s_3) \end{aligned} \quad (1.21.14)$$

These two A_1 molecular orbitals Ψ_1 and Ψ_2 , one bonding and one antibonding, are shown below.



The remaining two SALCs arising from the s orbitals of NH_3 :

$$\phi_3 = \frac{1}{\sqrt{6}}(2s_1 - s_2 - s_3) \quad (1.21.15)$$

and

$$\phi_4 = \frac{1}{\sqrt{2}}(s_2 - s_3) \quad (1.21.16)$$

form an orthogonal pair of molecular orbitals of E symmetry. We can show this by solving the secular determinant to find the orbital energies. The secular equations in this case are:

$$\begin{aligned} c_1(H_{33} - ES_{33}) + c_2(H_{34} - ES_{34}) &= 0 \\ c_1(H_{34} - ES_{34}) + c_2(H_{44} - ES_{44}) &= 0 \end{aligned} \quad (1.21.17)$$

Solving the secular determinant gives

$$E_{\pm} = \frac{-(2H_{34}S_{34} - H_{33} - H_{44}) \pm \sqrt{(2H_{34}S_{34} - H_{33} - H_{44})^2 - 4(1 - S_{34}^2)(H_{33}H_{44} - H_{34}^2)}}{2(1 - S_{34}^2)} \quad (1.21.18)$$


The integrals required are

$$\begin{aligned} H_{33} &= -9.2892 \text{ eV} \\ H_{44} &= -9.2892 \text{ eV} \\ H_{34} &= 0 \\ S_{34} &= 0 \end{aligned} \quad (1.21.19)$$

Using the fact that $H_{34} = S_{34} = 0$, the expression for the energies reduces to

$$E_{\pm} = \frac{(H_{33} + H_{44}) \pm (H_{33} - H_{44})}{2} \quad (1.21.20)$$

giving $E_+ = H_{33} = -9.2892 \text{ eV}$ and $E_- = H_{44} = -9.2892 \text{ eV}$. Each SALC therefore forms a molecular orbital by itself, and the two orbitals have the same energy; the two SALCs form an orthogonal pair of degenerate orbitals. These two molecular orbitals of E symmetry are shown below.



This page titled [1.21: Solving the Secular Equations](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.22: Summary of the Steps Involved in Constructing Molecular Orbitals

1. Choose a basis set of functions f_i consisting of the valence atomic orbitals on each atom in the system, or some chosen subset of these orbitals.
2. With the help of the appropriate character table, determine which irreducible representations are spanned by the basis set using Equation (15.20) to determine the number of times a_k that the k^{th} irreducible representation appears in the representation.

$$a_k = \frac{1}{h} \sum_C n_C \chi(g) \chi_k(g) \quad (1.22.1)$$

3. Construct the SALCs ϕ_i that transform as each irreducible representation using Equation 16.1

$$\phi_i = \sum_g \chi_k(g) g f_i \quad (1.22.2)$$


4. Write down expressions for the molecular orbitals by taking linear combinations of all the irreducible representations of the same symmetry species.
5. Write down the secular equations for the system.
6. Solve the secular determinant to obtain the energies of the molecular orbitals.
7. Substitute each energy in turn back into the secular equations and solve to obtain the coefficients appearing in your molecular orbital expressions in step 4.
8. Normalize the orbitals.

This page titled [1.22: Summary of the Steps Involved in Constructing Molecular Orbitals](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.23: A more complicated bonding example

As another example, we will use group theory to construct the molecular orbitals of H_2O (point group C_{2v}) using a basis set consisting of all the valence orbitals. The valence orbitals are a $1s$ orbital on each hydrogen, which we will label s_H and s'_H , and a $2s$ and three $2p$ orbitals on the oxygen, which we will label s_O, p_x, p_y, p_z giving a complete basis $(s_H, s'_H, s_O, p_x, p_y, p_z)$.

The first thing to do is to determine how each orbital transforms under the symmetry operations of the C_{2v} point group (E, C_2, σ_v and σ'_v), construct a matrix representation and determine the characters of each operation. The symmetry operations and axis system we will be using are shown below.



The orbitals transform in the following way

$$\begin{array}{ll}
 E & (s_H, s'_H, s_O, p_x, p_y, p_z) \rightarrow (s_H, s'_H, s_O, p_x, p_y, p_z) \\
 C_2 & (s_H, s'_H, s_O, p_x, p_y, p_z) \rightarrow (s'_H, s_H, s_O, -p_x, -p_y, p_z) \\
 \sigma_v(xz) & (s_H, s'_H, s_O, p_x, p_y, p_z) \rightarrow (s_H, s'_H, s_O, p_x, -p_y, p_z) \\
 \sigma'_v(yz) & (s_H, s'_H, s_O, p_x, p_y, p_z) \rightarrow (s'_H, s_H, s_O, -p_x, p_y, p_z)
 \end{array} \tag{1.23.1}$$

A short aside on constructing matrix representatives

After a little practice, you will probably be able to write matrix representatives straight away just by looking at the effect of the symmetry operations on the basis. However, if you are struggling a little the following procedure might help.

Remember that the matrix representatives are just the matrices we would have to multiply the left hand side of the above equations by to give the right hand side. In most cases they are very easy to work out. Probably the most straightforward way to think about it is that each column of the matrix shows where one of the original basis functions ends up. For example, the first column transforms the basis function s_H to its new position. The first column of the matrix can be found by taking the result on the right hand side of the above expressions, replacing every function that isn't s_H with a zero, putting the coefficient of s_H (1 or -1 in this example) in the position at which it occurs, and taking the transpose to give a column vector.

✓ Rotation

Consider the representative for the C_2 operation. The original basis $(s_H, s'_H, s_O, p_x, p_y, p_z)$ transforms into $(s'_H, s_H, s_O, -p_x, -p_y, p_z)$. The first column of the matrix therefore transforms s_H into s'_H . Taking the result and replacing all the other functions with zeroes gives $(0, s_H, 0, 0, 0, 0)$. The coefficient of s_H is 1, so the first column of the C_2 matrix representative is

$$\begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \tag{1.23.2}$$

Matrix representation, characters and SALCs

The matrix representatives and their characters are

$$\begin{array}{cccc}
 E & C_2 & \sigma_v & \sigma'_v \\
 \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} & \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \\
 \chi(E) = 6 & \chi(C_2) = 0 & \chi(\sigma_v) = 4 & \chi(\sigma'_v) = 2
 \end{array} \quad (1.23.3)$$

Now we are ready to work out which irreducible representations are spanned by the basis we have chosen. The character table for C_{2v} is:

C_{2v}	E	C_2	σ_v	σ'_v	$h = 4$
A_1	1	1	1	1	z, x^2, y^2, z^2
A_2	1	1	-1	-1	xy, R_z
B_1	1	-1	1	-1	x, xz, R_y
B_2	1	-1	-1	1	y, yz, R_x

As before, we use Equation (15.20) to find out the number of times each irreducible representation appears.

$$a_k = \frac{1}{h} \sum_C n_C \chi(g) \chi_k(g) \quad (1.23.5)$$

We have

$$\begin{aligned}
 a(A_1) &= \frac{1}{4}(1 \times 6 \times 1 + 1 \times 0 \times 1 + 1 \times 4 \times 1 + 1 \times 2 \times 1) &= 3 \\
 a(A_2) &= \frac{1}{4}(1 \times 6 \times 1 + 1 \times 0 \times 1 + 1 \times 4 \times -1 + 1 \times 2 \times -1) &= 0 \\
 a(B_1) &= \frac{1}{4}(1 \times 6 \times 1 + 1 \times 0 \times -1 + 1 \times 4 \times 1 + 1 \times 2 \times -1) &= 2 \\
 a(B_2) &= \frac{1}{4}(1 \times 6 \times 1 + 1 \times 0 \times -1 + 1 \times 4 \times -1 + 1 \times 2 \times 1) &= 1
 \end{aligned} \quad (1.23.6)$$

so the basis spans $3A_1 + 2B_1 + B_2$. Now we use the projection operators applied to each basis function f_i in turn to determine the SALCs $\phi_i = \sum_g \chi_k(g) g f_i$

The SALCs of A_1 symmetry are:

$$\begin{aligned}
 \phi(s_H) &= s_H + s'_H + s_H + s'_H &= 2(s_H + s'_H) \\
 \phi(s'_H) &= s'_H + s_H + s'_H + s_H &= 2(s_H + s'_H) \\
 \phi(s_O) &= s_O + s_O + s_O + s_O &= 4s_O \\
 \phi(p_x) &= p_x - p_x + p_x - p_x &= 0 \\
 \phi(p_y) &= p_y - p_y + p_y - p_y &= 0 \\
 \phi(p_z) &= p_z + p_z + p_z + p_z &= 4p_z
 \end{aligned} \quad (1.23.7)$$

The SALCs of B_1 symmetry are:

$$\begin{aligned}
 \phi(s_H) &= s_H - s'_H + s_H - s'_H &= 2(s_H - s'_H) \\
 \phi(s'_H) &= s'_H - s_H + s'_H - s_H &= 2(s'_H - s_H) \\
 \phi(s_O) &= s_O - s_O + s_O - s_O &= 0 \\
 \phi(p_x) &= p_x + p_x + p_x + p_x &= 4p_x \\
 \phi(p_y) &= p_y + p_y - p_y - p_y &= 0 \\
 \phi(p_z) &= p_z - p_z + p_z - p_z &= 0
 \end{aligned} \quad (1.23.8)$$

The SALCs of B_2 symmetry are:

$$\begin{aligned}
 \phi(s_H) &= s_H - s'_H - s_H + s'_H = 0 \\
 \phi(s'_H) &= s'_H - s_H - s'_H + s_H = 0 \\
 \phi(s_O) &= s_O - s_O - s_O + s_O = 0 \\
 \phi(p_x) &= p_x + p_x - p_x - p_x = 0 \\
 \phi(p_y) &= p_y + p_y + p_y + p_y = 4p_y \\
 \phi(p_z) &= p_z - p_z - p_z + p_z = 0
 \end{aligned}
 \tag{1.23.9}$$

After normalization, our SALCs are therefore:

A_1 symmetry

$$\begin{aligned}
 \phi_1 &= \frac{1}{\sqrt{2}}(s_H + s'_H) \\
 \phi_2 &= s_O \\
 \phi_3 &= p_z
 \end{aligned}
 \tag{1.23.10}$$

B_1 symmetry

$$\begin{aligned}
 \phi_4 &= \frac{1}{\sqrt{2}}(s_H - s'_H) \\
 \phi_5 &= p_x
 \end{aligned}
 \tag{1.23.11}$$

B_2 symmetry

$$\phi_6 = p_y
 \tag{1.23.12}$$

Note that we only take one of the first two SALCs generated by the B_1 projection operator since one is a simple multiple of the other (i.e. they are not linearly independent). We can therefore construct three molecular orbitals of A_1 symmetry, with the general form

$$\begin{aligned}
 \Psi(A_1) &= c_1\phi_1 + c_2\phi_2 + c_3\phi_3 \\
 &= c'_1(s_H + s'_H) + c_2s_O + c_3p_z \quad \text{where } c'_1 = \frac{c_1}{\sqrt{2}}
 \end{aligned}
 \tag{1.23.13}$$


two molecular orbitals of B_1 symmetry, of the form

$$\begin{aligned}
 \Psi(B_1) &= c_4\phi_4 + c_5\phi_5 \\
 &= c'_4(s_H - s'_H) + c_5p_x
 \end{aligned}
 \tag{1.23.14}$$

and one molecular orbital of B_2 symmetry

$$\begin{aligned}
 \Psi(B_2) &= \phi_6 \\
 &= p_y
 \end{aligned}
 \tag{1.23.15}$$

To work out the coefficients $c_1 - c_5$ and determine the orbital energies, we would have to solve the secular equations for each set of orbitals in turn. We are not dealing with a conjugated p system, so in this case Hückel theory cannot be used and the various H_{ij} and S_{ij} integrals would have to be calculated numerically and substituted into the secular equations. This involves a lot of tedious algebra, which we will leave out for the moment. The LCAO orbitals determined above are an approximation of the true molecular orbitals of water, which are shown on the right. As we have shown using group theory, the A_1 molecular orbitals involve the oxygen $2s$ and $2p_z$ atomic orbitals and the sum $s_H + s'_H$ of the hydrogen $1s$ orbitals. The B_1 molecular orbitals involve the oxygen $2p_x$ orbital and the difference $s_H - s'_H$ of the two hydrogen $1s$ orbitals, and the B_2 molecular orbital is essentially an oxygen $2p_y$ atomic orbital.




Contributors and Attributions

[Claire Vallance](#) (University of Oxford)

This page titled [1.23: A more complicated bonding example](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.24: Molecular Vibrations

Vibrational motion in diatomic molecules is often discussed within the context of the simple harmonic oscillator in quantum mechanics. A diatomic molecule has only a single bond that can vibrate; we say it has a single *vibrational mode*. As you may expect, the vibrational motions of polyatomic molecules are much more complicated than those in a diatomic. Firstly, there are more bonds that can vibrate; and secondly, in addition to stretching vibrations, the only type of vibration possible in a diatomic, we can also have bending and torsional vibrational modes. Since changing one bond length in a polyatomic will often affect the length of nearby bonds, we cannot consider the vibrational motion of each bond in isolation; instead we talk of *normal modes* involving the concerted motion of groups of bonds. As a simple example, the normal modes of a linear triatomic molecule are shown below.



Once we know the symmetry of a molecule at its equilibrium structure, group theory allows us to predict the vibrational motions it will undergo using exactly the same tools we used above to investigate molecular orbitals. Each vibrational mode transforms as one of the irreducible representations of the molecule's point group. Before moving on to an example, we will quickly review how to determine the number of vibrational modes in a molecule.

Molecular degrees of freedom – determining the number of normal vibrational modes

An atom can undergo only translational motion, and therefore has three degrees of freedom corresponding to motion along the x , y , and z Cartesian axes. Translational motion in any arbitrary direction can always be expressed in terms of components along these three axes. When atoms combine to form molecules, each atom still has three degrees of freedom, so the molecule as a whole has $3N$ degrees of freedom, where N is the number of atoms in the molecule. However, the fact that each atom in a molecule is bonded to one or more neighboring atoms severely hinders its translational motion, and also ties its motion to that of the atoms to which it is attached. For these reasons, while it is entirely possible to describe molecular motions in terms of the translational motions of individual atoms (we will come back to this in the next section), we are often more interested in the motions of the molecule as a whole. These may be divided into three types: translational; rotational and vibrational.

Just as for an individual atom, the molecule as a whole has three degrees of translational freedom, leaving $3N - 3$ degrees of freedom in rotation and vibration.

The number of rotational degrees of freedom depends on the structure of the molecule. In general, there are three possible rotational degrees of freedom, corresponding to rotation about the x , y , and z Cartesian axes. A non-linear polyatomic molecule does indeed have three rotational degrees of freedom, leaving $3N - 6$ degrees of freedom in vibration (i.e. $3N - 6$ vibrational modes). In a linear molecule, the situation is a little different. It is generally accepted that to be classified as a true rotation, a motion must change the position of one or more of the atoms. If we define the z axis as the molecular axis, we see that spinning the molecule about the axis does not move any of the atoms from their original position, so this motion is not truly a rotation. Consequently, a linear molecule has only two degrees of rotational freedom, corresponding to rotations about the x and y axis. This type of molecule has $3N - 5$ degrees of freedom left for vibration, or $3N - 5$ vibrational modes.

In summary:

- A linear molecule has $3N - 5$ vibrational modes
- A non-linear molecule has $3N - 6$ vibrational modes.


Determining the Symmetries of Molecular Motions

We mentioned above that the procedure for determining the normal vibrational modes of a polyatomic molecule is very similar to that used in previous sections to construct molecular orbitals. In fact, virtually the only difference between these two applications of group theory is the choice of basis set.

As we have already established, the motions of a molecule may be described in terms of the motions of each atom along the x , y and z axis. Consequently, it probably won't come as too much of a surprise to discover that a very useful basis for describing molecular motions comprises a set of (x, y, z) axes centered on each atom. This basis is usually known as the $3N$ Cartesian basis (since there are $3N$ Cartesian axes, 3 axes for each of the N atoms in the molecule). Note that each molecule will have a different $3N$ Cartesian basis, just as every molecule has a different atomic orbital basis.

Our first task in investigating motions of a particular molecule is to determine the characters of the matrix representatives for the $3N$ Cartesian basis under each of the symmetry operations in the molecular point group. We will use the H_2O molecule, which has C_{2v} symmetry, as an example.

H_2O has three atoms, so the $3N$ Cartesian basis will have 9 elements. The basis vectors are shown in the diagram below.



One way of determining the characters would be to construct all of the matrix representatives and take their traces. While you are more than welcome to try this approach if you want some practice at constructing matrix representatives, there is an easier way. Recall that we can also determine the character of a matrix representative under a particular symmetry operation by stepping through the basis functions and applying the following rules:

- i. Add 1 to the character if the basis function is unchanged by the symmetry operation;
- ii. Add -1 to the character if the basis function changes sign under the symmetry operation;
- iii. Add 0 to the character if the basis function moves when the symmetry operation is applied.

For H_2O , this gives us the following characters for the $3N$ Cartesian basis (check that you can obtain this result using the rules above and the basis vectors as drawn in the figure):

$$\begin{array}{l} \text{Operation: } E \quad C_2 \quad \sigma_v(xz) \quad \sigma'_v(yz) \\ \chi_{3N} : \quad \quad 9 \quad -1 \quad 3 \quad 1 \end{array} \quad (24.1)$$

There is an even quicker way to work out the characters of the $3N$ Cartesian basis if you have a character table in front of you. The character for the *Cartesian* basis is simply the sum of the characters for the x , y , and z (or T_x , T_y , and T_z) functions listed in the character table. To get the character for the $3N$ Cartesian basis, simply multiply this by the number of atoms in the molecule that are unshifted by the symmetry operation.

The C_{2v} character table is shown below.

C_{2v}	E	C_2	σ_v	σ'_v	$h = 4$
A_1	1	1	1	1	z, x^2, y^2, z^2
A_2	1	1	-1	-1	xy, R_z
B_1	1	-1	1	-1	x, xz, R_y
B_2	1	-1	-1	1	y, yz, R_x

(24.2)

x transforms as B_1 , y as B_2 , and z as A_1 , so the characters for the Cartesian basis are

$$\begin{array}{l} \text{Operation: } E \quad C_2 \quad \sigma_v(xz) \quad \sigma'_v(yz) \\ \chi_{3N} : \quad \quad 3 \quad -1 \quad 1 \quad 1 \end{array} \quad (24.3)$$

We multiply each of these by the number of unshifted atoms (3 for the identity operation, 1 for C_2 , 3 for σ_v and 1 for σ'_v) to obtain the characters for the $3N$ Cartesian basis.

$$\chi_{3N} : \quad 9 \quad -1 \quad 3 \quad 1 \quad (24.4)$$

Reassuringly, we obtain the same characters as we did previously. Which of the three methods you use to get to this point is up to you.


We now have the characters for the molecular motions (described by the $3N$ Cartesian basis) under each symmetry operation. At this point, we want to separate these characters into contributions from translation, rotation, and vibration. This turns out to be a very straightforward task. We can read the characters for the translational and rotational modes directly from the character table, and we obtain the characters for the vibrations simply by subtracting these from the $3N$ Cartesian characters we've just determined. The characters for the translations are the same as those for χ_{Cart} . We find the characters for the rotations by adding together the characters for R_x , R_y , and R_z from the character table (or just R_x and R_y if the molecule is linear). For H_2O , we have:

Operation:	E	C_2	$\sigma_v(xz)$	$\sigma'_v(yz)$	
χ_{3N} :	9	-1	3	1	
χ_{Trans} :	3	-1	1	1	(24.5)
χ_{Rot} :	3	-1	-1	-1	
$\chi_{\text{Vib}} = \chi_{3N} - \chi_{\text{Trans}} - \chi_{\text{Rot}}$:	3	1	3	1	

The characters in the final row are the sums of the characters for all of the molecular vibrations. We can find out the symmetries of the individual vibrations by using the reduction equation (Equation (15.20)) to determine the contribution from each irreducible representation.

In many cases you won't even need to use the equation, and can work out which irreducible representations are contributing just by inspection of the character table. In the present case, the only combination of irreducible representations that can give the required values for χ_{Vib} is $2A_1 + B_1$. As an exercise, you should make sure you are also able to obtain this result using the reduction equation.

So far this may all seem a little abstract, and you probably want to know what the vibrations of H_2O actually look like. For a molecule with only three atoms, it is fairly easy to identify the possible vibrational modes and to assign them to the appropriate irreducible representation.



For a larger molecule, the problem may become much more complex, and in that case we can generate the SALCs of the $3N$ Cartesian basis, which will tell us the atomic displacements associated with each vibrational mode. We will do this now for H_2O .

Atomic displacements using the $3N$ Cartesian basis

As before, we generate the SALCs of each symmetry by applying the appropriate projection operator to each of the basis functions (or in this case, basis vectors) f_i in turn.

$$\phi_i = \sum_g \chi_k(g) g f_i \quad (24.6)$$

In this case we have 9 basis vectors, which we will label $x_H, y_H, z_H, x_O, y_O, z_O, x_{H'}, y_{H'}, z_{H'}$, describing the displacements of the two H atoms and the O atom along Cartesian axes. For the SALCs of A_1 symmetry, applying the projection operator to each basis vector in turn gives (check that you can obtain this result):

$$\begin{aligned}
 \phi_1(x_H) &= x_H - x_{H'} + x_H - x_{H'} &= 2x_H - 2x_{H'} \\
 \phi_2(y_H) &= y_H - y_{H'} - y_H + y_{H'} &= 0 \\
 \phi_3(z_H) &= z_H + z_{H'} + z_H + z_{H'} &= 2z_H + 2z_{H'} \\
 \phi_4(x_O) &= x_O - x_O + x_O - x_O &= 0 \\
 \phi_5(y_O) &= y_O - y_O - y_O + y_O &= 0 \\
 \phi_6(z_O) &= z_O + z_O + z_O + z_O &= 4z_O \\
 \phi_7(x_{H'}) &= x_{H'} - x_H + x_{H'} - x_H &= 2x_{H'} - 2x_H \\
 \phi_8(y_{H'}) &= y_{H'} - y_H - y_{H'} + y_H &= 0 \\
 \phi_9(z_{H'}) &= z_{H'} + z_H + z_{H'} + z_H &= 2z_{H'} + 2z_H
 \end{aligned} \quad (24.7)$$

We see that the motion characteristic of an A_1 vibration (which we have identified as the symmetric stretch and the bending vibration) may be summarized as follows:

- i. $2(x_H - x_{H'})$ - the two hydrogen atoms move in opposite directions along the x axis.
- ii. $2(z_H + z_{H'})$ - the two hydrogen atoms move in the same direction along the z axis.
- iii. $4z_O$ - the oxygen atom moves along the z axis.
- iv. There is no motion of any of the atoms in the y direction.

The asymmetric stretch has B_1 symmetry, and applying the projection operator in this case gives:

$$\begin{aligned}
 \phi_1(x_H) &= x_H + x_{H'} + x_H + x_{H'} = 2x_H + 2x_{H'} \\
 \phi_2(y_H) &= y_H + y_{H'} - y_H - y_{H'} = 0 \\
 \phi_3(z_H) &= z_H - z_{H'} + z_H - z_{H'} = 2z_H - 2z_{H'} \\
 \phi_4(x_O) &= x_O + x_O + x_O + x_O = 4x_O \\
 \phi_5(y_O) &= y_O + y_O - y_O - y_O = 0 \\
 \phi_6(z_O) &= z_O - z_O + z_O - z_O = 0 \\
 \phi_7(x_{H'}) &= x_{H'} + x_H + x_{H'} + x_H = 2x_{H'} + 2x_H \\
 \phi_8(y_{H'}) &= y_{H'} + y_H - y_{H'} - y_H = 0 \\
 \phi(z_{H'}) &= z_{H'} - z_H + z_{H'} - z_H = 2z_{H'} - 2z_H
 \end{aligned}
 \tag{24.8}$$

In this vibrational mode, the two H atoms move in the same direction along the x axis and in opposite directions along the z axis.

We have now shown how group theory may be used together with the $3N$ Cartesian basis to identify the symmetries of the translational, rotational and vibrational modes of motion of a molecule, and also to determine the atomic displacements associated with each vibrational mode.

Molecular vibrations using internal coordinates

While it was fairly straightforward to investigate the atomic displacements associated with each vibrational mode of H_2O using the $3N$ Cartesian basis, this procedure becomes more complicated for larger molecules. Also, we are often more interested in how bond lengths and angles change in a vibration, rather than in the Cartesian displacements of the individual atoms. If we are only interested in looking at molecular *vibrations*, we can use a different procedure from that described above, and start from a basis of *internal coordinates*. Internal coordinates are simply a set of bond lengths and bond angles, which we can use as a basis for generating representations and, eventually, SALCs. Since bond lengths and angles do not change during translational or rotational motion, no information will be obtained on these types of motion.

For H_2O , the three internal coordinates of interest are the two OH bond lengths, which we will label r and r' , and the HOH bond angle, which we will label θ . If we wanted to, we could separate our basis into two different bases, one consisting only of bond lengths, to describe stretching vibrations, and one consisting of only bond angles, to describe bending vibrations. However, the current example is simple enough to treat all the basis functions together.

As usual, our first step is to work out the characters of the matrix representatives for this basis under each symmetry operation. The effects of the various transformations on our chosen basis, and the characters of the corresponding representatives, are:

$$\begin{aligned}
 E(r, r', \theta) &= (r, r', \theta) & \chi(E) &= 3 \\
 C_2(r, r', \theta) &= (r', r, \theta) & \chi(C_2) &= 1 \\
 \sigma_v(xz)(r, r', \theta) &= (r, r', \theta) & \chi(\sigma_v) &= 3 \\
 \sigma'_v(yz)(r, r', \theta) &= (r', r, \theta) & \chi(\sigma'_v) &= 1
 \end{aligned}
 \tag{24.9}$$

These are the same characters as we found before using the $3N$ Cartesian basis, and as before, we can see by inspection of the character table that the representation may be reduced down to the sum of irreducible representations $2A_1 + B_1$. We can now work out the symmetry adapted linear combinations of our new basis set to see how the bond lengths and angle change as H_2O vibrates in each of the three vibrational modes.

Again, we will use the projection operator $\phi_i = \sum_g \chi_k(g) g f_i$ applied to each basis function in turn.

Firstly, the A_1 vibrations:

$$\begin{aligned}
 \phi_1(r) &= r + r' + r + r' = 2(r + r') \\
 \phi_2(r') &= r' + r + r' + r = 2(r' + r) \\
 \phi_3(\theta) &= \theta + \theta + \theta + \theta = 4\theta
 \end{aligned}
 \tag{24.10}$$

From these SALCs, we can identify ϕ_1 (and ϕ_2 , which is identical) with the symmetric stretch, in which both bond lengths change in phase with each other, and ϕ_3 with the bend.

Now for the B_1 vibration:

$$\begin{aligned}\phi_4(r) &= r - r' + r - r' &= 2(r - r') \\ \phi_5(r') &= r' - r + r' - r &= 2(r' - r) \\ \phi_6(\theta) &= \theta - \theta + \theta - \theta &= 0\end{aligned}\tag{24.11}$$

ϕ_4 and ϕ_5 are not linearly independent, and either one may be chosen to describe the asymmetric stretch, in which one bond lengthens as the other shortens.

Note: When using internal coordinates, it is important that all of the coordinates in the basis are linearly independent. If this is the case then the number of internal coordinates in the basis will be the same as the number of vibrational modes ($3N - 5$ or $3N - 6$, depending on whether the molecule is linear or non-linear). This requirement is satisfied in the H_2O example above. For a less straightforward example, consider the methane molecule, CH_4 . It might appear that we could choose a basis made up of the four $C-H$ bond lengths and the six $H-C-H$ bond angles. However, this would give us 10 basis functions, and CH_4 has only 9 vibrational modes. This is due to the fact that the bond angles are not all independent of each other. It can be tricky to come up with the appropriate internal coordinate basis to describe all of the molecular motions, but all is not lost. Even if you can't work out the appropriate bond angles to choose, you can always take a basis of bond lengths to investigate the stretching vibrations of a molecule. If you want to know the symmetries of the bending vibrations, you can use the $3N$ Cartesian basis method to determine the symmetries of all of the vibrational modes and compare these with the stretching mode symmetries to identify the bending modes.

Contributors and Attributions

Claire Vallance (University of Oxford)

This page titled [1.24: Molecular Vibrations](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.25: Summary of applying group theory to molecular motions

1. Atomic or molecular translations transform in the same way as the x, y, z (or T_x, T_y, T_z) functions listed in the character tables.
2. Molecular rotations transform in the same way as the R_x, R_y, R_z functions listed in the character tables.
3. The irreducible representations spanned by the motions of a polyatomic molecule may be determined using the $3N$ Cartesian basis, made up of x, y, z axes on each atom. The characters of the matrix representatives are best determined using a table as follows:

Operation:	List the symmetry operations in the point group	
Γ_{Cart}	List the characters for $x + y + z$ (from the character table) for each operation	(1.25.1)
$N_{\text{unshifted}}$	List the number of atoms in the molecule that are unshifted by each symmetry operation	
Γ_{3N}	Take the product of the previous two rows to give the characters for Γ_{3N}	

4. The irreducible representations spanned by the molecular vibrations are determined by first subtracting the characters for rotations and translations from the characters for Γ_{3N} to give the characters for Γ_{vib} and then using the reduction formula or inspection of the character table to identify the irreducible representations contributing to Γ_{vib} .
5. The molecular displacements for the vibrations of each symmetry may be determined by using projection operators on the $3N$ Cartesian basis vectors to generate SALCs.
6. Alternatively, a basis of internal coordinates (bond lengths and angles) may be used to investigate stretching and bending vibrations. Determine the characters, identify the irreducible representations, and construct SALCs.

Contributors and Attributions

[Claire Vallance](#) (University of Oxford)

This page titled [1.25: Summary of applying group theory to molecular motions](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.26: Group theory and Molecular Electronic States

Firstly, it is important that you understand the difference between a *molecular orbital* and an *electronic state*.

A strict definition of a molecular orbital is that it is a ‘one electron wavefunction’, i.e. a solution to the Schrödinger equation for the molecule. A complete one electron wavefunction (orbital) is a product of a spatial function, describing the orbital angular momentum and ‘shape’ of the orbital, and a spin function, describing the spin angular momentum.

$$\Psi = \Psi_{\text{spatial}}\Psi_{\text{spin}} \quad (26.1)$$

In common usage, the word ‘orbital’ is often used to refer only to the spatial part of the ‘true’ orbital. For example, in atoms we generally talk about ‘*s* orbitals’ or ‘*p* orbitals’ rather than ‘*s* spatial wavefunctions’ and ‘*p* spatial wavefunctions’. In this context, two electrons with opposite spins may occupy one spatial orbital. A more rigorous way of saying this would be to state that a given spatial wavefunction may be paired with two different spin wavefunctions (one corresponding to a ‘spin up’ electron and one to a ‘spin down’ electron).

An electronic state is defined by the electron configuration of the system, and by the quantum numbers of each electron contributing to that configuration. Each electronic state corresponds to one of the energy levels of the molecule. These energy levels will obviously depend on the molecular orbitals that are occupied, and their energies, but they also depend on the way in which the electrons within the various molecular orbitals interact with each other. Interactions between the electrons are essentially determined by the relative orientations of the magnetic moments associated with their orbital and spin angular momenta, which is where the dependence on quantum numbers comes in. A given electron configuration will often give rise to a number of different electronic states if the electrons may be arranged in different ways (with different quantum numbers) within the occupied orbitals.

Last year you were introduced to the idea of atomic states, and learnt how to label the states arising from a given electron configuration using term symbols of the form $^{2S+1}L_J$. Term symbols of this form define the spin, orbital and total angular momenta of the state, which in turn determine its energy. Molecular states, containing contributions from a number of molecular orbitals, are more complicated. For example, a given molecular orbital will generally contain contributions from several different atomic orbitals, and as a result, electrons cannot easily be assigned an *l* quantum number. Instead of using term symbols, molecular states are usually labeled according to their symmetry (the exception to this is linear molecules, for which conventional term symbols may still be used, albeit with a few modifications from the atomic case).

We can determine the symmetry of an electronic state by taking the direct product of the irreducible representations for all of the electrons involved in that state (the irreducible representation for each electron is simply the irreducible representation for the molecular orbital that it occupies). Usually we need only consider unpaired electrons. Closed shell species, in which all electrons are paired, almost always belong to the totally symmetric irreducible representation in the point group of the molecule.

An example is the molecular orbitals of butadiene, which belongs to the C_{2h} point group. Since all electrons are paired, the overall symmetry of the state is A_g , and the label for the state once the spin multiplicity is included is 1A_g . We could have arrived at the same result by taking the direct product of the irreducible representations for each electron. There are two electrons in orbitals with A_u symmetry, and two in orbitals with B_g symmetry, so overall we have:

$$A_u \otimes A_u \otimes B_g \otimes B_g = A_g \quad (26.2)$$

This page titled [1.26: Group theory and Molecular Electronic States](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.27: Spectroscopy - Interaction of Atoms and Molecules with Light

In our final application of group theory, we will investigate the way in which symmetry considerations influence the interaction of light with matter. We have already used group theory to learn about the molecular orbitals in a molecule. In this section we will show that it may also be used to predict which electronic states may be accessed by absorption of a photon. We may also use group theory to investigate how light may be used to excite the various vibrational modes of a polyatomic molecule.

Last year, you were introduced to spectroscopy in the context of electronic transitions in atoms. You learned that a photon of the appropriate energy is able to excite an electronic transition in an atom, subject to the following selection rules:

$$\begin{aligned}
 \Delta n &= \text{integer} \\
 \Delta l &= \pm 1 \\
 \Delta L &= 0, \pm 1 \\
 \Delta S &= 0 \\
 \Delta J &= 0, \pm 1; J = 0 \leftrightarrow J = 0
 \end{aligned}
 \tag{27.1}$$

What you may not have learned is where these selection rules come from. In general, different types of spectroscopic transition obey different selection rules. The transitions you have come across so far involve changing the *electronic* state of an atom, and involve absorption of a photon in the UV or visible part of the electromagnetic spectrum. There are analogous electronic transitions in molecules, which we will consider in more detail shortly. Absorption of a photon in the infrared (IR) region of the spectrum leads to *vibrational* excitation in molecules, while photons in the microwave (MW) region produce *rotational* excitation. Each type of excitation obeys its own selection rules, but the general procedure for determining the selection rules is the same in all cases. It is simply to determine the conditions under which the probability of a transition is not identically zero.

The first step in understanding the origins of selection rules must therefore be to learn how transition probabilities are calculated. This requires some quantum mechanics.

Last year, you learned about operators, eigenvalues and eigenfunctions in quantum mechanics. You know that if a function is an eigenfunction of a particular operator, then operating on the eigenfunction with the operator will return the observable associated with that state, known as the eigenvalue (i.e. $\hat{A}\Psi = a\Psi$). What you may not know is that operating on a function that is NOT an eigenfunction of the operator leads to a change in state of the system. In the transitions we will be considering, the molecule interacts with the *electric* field of the light (as opposed to NMR spectroscopy, in which the nuclei interact with the magnetic field of the electromagnetic radiation). These transitions are called *electric dipole transitions*, and the operator we are interested in is the *electric dipole operator*, usually given the symbol $\hat{\mu}$, which describes the electric field of the light.

If we start in some initial state Ψ_i , operating on this state with $\hat{\mu}$ gives a new state, $\hat{\mu}\Psi_i$. If we want to know the probability of ending up in some particular final state Ψ_f , the probability amplitude is simply given by the overlap integral between $\hat{\mu}\Psi_i$ and Ψ_f . This probability amplitude is called the *transition dipole moment*, and is given the symbol μ_{fi} .

$$\mu_{fi} = \langle \Psi_f | \hat{\mu} | \Psi_i \rangle
 \tag{27.2}$$

Physically, the transition dipole moment may be thought of as describing the ‘kick’ the electron receives or imparts to the electric field of the light as it undergoes a transition. The transition probability is given by the square of the probability amplitude.

$$P_{fi} = \mu_{fi}^2 = |\langle \Psi_f | \hat{\mu} | \Psi_i \rangle|^2
 \tag{27.3}$$

Hopefully it is clear that in order to determine the selection rules for an electric dipole transition between states Ψ_i and Ψ_f , we need to find the conditions under which μ_{fi} can be non-zero. One way of doing this would be to write out the equations for the two wavefunctions (which are functions of the quantum numbers that define the two states) and the electric dipole moment operator, and just churn through the integrals. By examining the result, it would then be possible to decide what restrictions must be imposed on the quantum numbers of the initial and final states in order for a transition to be allowed, leading to selection rules of the type listed above for atoms. However, many selection rules may be derived with a lot less work, based simply on symmetry considerations.

In section 17, we showed how to use group theory to determine whether or not an integral may be non-zero. This forms the basis of our consideration of selection rules.

Electronic transitions in molecules


Assume that we have a molecule in some initial state Ψ_i . We want to determine which final states Ψ_f can be accessed by absorption of a photon. Recall that for an integral to be non-zero, the representation for the integrand must contain the totally symmetric irreducible representation. The integral we want to evaluate is

$$\hat{\mu}_{fi} = \int \Psi_f^* \hat{\mu} \Psi_i d\tau \quad (27.4)$$

so we need to determine the symmetry of the function $\Psi_f^* \hat{\mu} \Psi_i$. As we learned in Section 18, the product of two functions transforms as the direct product of their symmetry species, so all we need to do to see if a transition between two chosen states is allowed is work out the symmetry species of Ψ_f , $\hat{\mu}$ and Ψ_i , take their direct product, and see if it contains the totally symmetric irreducible representation for the point group of interest. Equivalently (as explained in Section 18), we can take the direct product of the irreducible representations for $\hat{\mu}$ and Ψ_i and see if it contains the irreducible representation for Ψ_f . This is best illustrated using a couple of examples.


Earlier in the course, we learned how to determine the symmetry molecular orbitals. The symmetry of an electronic state is found by identifying any unpaired electrons and taking the direct product of the irreducible representations of the molecular orbitals in which they are located. The ground state of a closed-shell molecule, in which all electrons are paired, always belongs to the totally symmetric irreducible representation⁷. As an example, the electronic ground state of NH_3 , which belongs to the C_{3v} point group, has A_1 symmetry. To find out which electronic states may be accessed by absorption of a photon, we need to determine the irreducible representations for the electric dipole operator $\hat{\mu}$. Light that is linearly polarized along the x , y , and z axes transforms in the same way as the functions x , y , and z in the character table⁸. From the C_{3v} character table, we see that x - and y -polarized light transforms as E , while z -polarized light transforms as A_1 . Therefore:

- For x - or y -polarized light, $\Gamma_{\hat{\mu}} \otimes \Gamma_{\Psi_1}$ transforms as $E \otimes A_1 = E$. This means that absorption of x - or y -polarized light by ground-state NH_3 (see figure below left) will excite the molecule to a state of E symmetry.
- For z -polarized light, $\Gamma_{\hat{\mu}} \otimes \Gamma_{\Psi_1}$ transforms as $A_1 \otimes A_1 = A_1$. Absorption of z -polarized light by ground state NH_3 (see figure below right) will excite the molecule to a state of A_1 symmetry.



Of course, the photons must also have the appropriate energy, in addition to having the correct polarization to induce a transition.

We can carry out the same analysis for H_2O , which belongs to the C_{2v} point group. We showed previously that H_2O has three molecular orbitals of A_1 symmetry, two of B_1 symmetry, and one of B_2 symmetry, with the ground state having A_1 symmetry. In the C_{2v} point group, x -polarized light has B_1 symmetry, and can therefore be used to excite electronic states of this symmetry; y -polarized light has B_2 symmetry, and may be used to access the B_2 excited state; and z -polarized light has A_1 symmetry, and may be used to access higher lying A_1 states. Consider our previous molecular orbital diagram for H_2O .




The electronic ground state has two electrons in a B_2 orbital, giving a state of A_1 symmetry ($B_2 \otimes B_2 = A_1$). The first excited electronic state has the configuration $(1B_2)^1(3A_1)^1$ and its symmetry is $B_2 \otimes A_1 = B_2$. It may be accessed from the ground state by a y -polarized photon. The second excited state is accessed from the ground state by exciting an electron to the $2B_1$ orbital. It has the configuration $(1B_2)^1(2B_1)^1$, its symmetry is $B_2 \otimes B_1 = A_2$. Since neither x -, y - or z -polarized light transforms as A_2 , this state may not be excited from the ground state by absorption of a single photon.

Vibrational transitions in molecules


Similar considerations apply for vibrational transitions. Light polarized along the x , y , and z axes of the molecule may be used to excite vibrations with the same symmetry as the x , y and z functions listed in the character table.

For example, in the C_{2v} point group, x -polarized light may be used to excite vibrations of B_1 symmetry, y -polarized light to excite vibrations of B_2 symmetry, and z -polarized light to excite vibrations of A_1 symmetry. In H_2O , we would use z -polarized light to excite the symmetric stretch and bending modes, and x -polarized light to excite the asymmetric stretch. Shining y -polarized light onto a molecule of H_2O would not excite any vibrational motion.



Raman Scattering

If there are vibrational modes in the molecule that may not be accessed using a single photon, it may still be possible to excite them using a two-photon process known as Raman scattering⁹. An energy level diagram for Raman scattering is shown below.



The first photon excites the molecule to some high-lying intermediate state, known as a *virtual state*. Virtual states are not true stationary states of the molecule (i.e. they are not eigenfunctions of the molecular Hamiltonian), but they can be thought of as stationary states of the 'photon + molecule' system. These types of states are extremely short lived, and will quickly emit a photon to return the system to a stable molecular state, which may be different from the original state. Since there are two photons (one absorbed and one emitted) involved in Raman scattering, which may have different polarizations, the transition dipole for a Raman transition transforms as one of the Cartesian products x^2 , y^2 , z^2 , xy , xz , yz listed in the character tables.

Vibrational modes that transform as one of the Cartesian products may be excited by a Raman transition, in much the same way as modes that transform as x , y , or z may be excited by a **one-photon** vibrational transition.

In H_2O , all of the vibrational modes are accessible by ordinary one-photon vibrational transitions. However, they may also be accessed by Raman transitions. The Cartesian products transform as follows in the C_{2v} point group.

$$\begin{array}{ll} A_1 & x^2, y^2, z^2 \\ A_2 & sy \end{array} \quad \begin{array}{ll} B_1 & xz \\ B_2 & yz \end{array} \quad (27.5)$$

The symmetric stretch and the bending vibration of water, both of A_1 symmetry, may therefore be excited by any Raman scattering process involving two photons of the same polarization (x -, y - or z -polarized). The asymmetric stretch, which has B_1 symmetry, may be excited in a Raman process in which one photon is x -polarized and the other z -polarized.

⁷ It is important not to confuse *molecular orbitals* (the energy levels that individual electrons may occupy within the molecule) with *electronic states* (arising from the different possible arrangements of *all* the molecular electrons amongst the molecular orbitals, e.g. the electronic states of NH_3 are NOT the same thing as the molecular orbitals we derived earlier in the course. These orbitals were an incomplete set, based only on the valence s electrons in the molecule. Inclusion of the p electrons is required for a full treatment of the electronic states. The H_2O example above should hopefully clarify this point.

⁸ ' x -polarized' means that the electric vector of the light (an electromagnetic wave) oscillates along the direction of the x axis.

⁹ You will cover Raman scattering (also known as Raman spectroscopy) in more detail in later courses. The aim here is really just to alert you to its existence and to show how it may be used to access otherwise inaccessible vibrational modes.

This page titled [1.27: Spectroscopy - Interaction of Atoms and Molecules with Light](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.28: Summary

Hopefully this course has given you a reasonable introduction to the qualitative description of molecular symmetry, and also to the way in which it can be used quantitatively within the context of group theory to predict important molecular properties. These main things you should have learnt in this text are:

1. How to identify the symmetry elements possessed by a molecule and assign it to a point group.
2. The consequences of symmetry for chirality and polarity of molecules.
3. The effect of applying two or more symmetry operations consecutively (group multiplication)
4. How to construct a matrix representation of a group, starting from a suitable set of basis functions.
5. How to determine the irreducible representations spanned by a basis set, and construct symmetry adapted linear combinations (SALCs) of the original basis functions that transform as the irreducible representations of the group.
6. How to construct molecular orbitals by taking linear combinations of SALCs of the same symmetry species.
7. How to set up and solve the secular equations for the molecule in order to find the molecular energy levels and orbital coefficients – “Extra for experts”, though you will cover this in later courses
8. How to determine the symmetries of the various modes of motion (translational, rotational and vibrational) of a polyatomic molecule, and the symmetries of individual vibrational modes.
9. How to determine the atomic displacements in a given vibrational mode by constructing SALCs in the $3N$ Cartesian basis.
10. How to determine atomic displacements in stretching and bending vibrations using internal coordinates.
11. The consequences of symmetry for the selection rules governing excitation to different electronic and vibrational states.

Contributors and Attributions

[Claire Vallance](#) (University of Oxford)

This page titled [1.28: Summary](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.29: Appendix A

Proof that the character of a matrix representative is invariant under a similarity transform

A property of traces of matrix products is that they are invariant under cyclic permutation of the matrices.

i.e. $\text{tr}[ABC] = \text{tr}[BCA] = \text{tr}[CAB]$. For the character of a matrix representative of a symmetry operation g , we therefore have:

$$\chi(g) = \text{tr}[\Gamma(g)] = \text{tr}[C\Gamma'(g)C^{-1}] = \text{tr}[\Gamma'(g)C^{-1}C] = \text{tr}[\Gamma'(g)] = \chi'(g) \quad (29.1)$$

The trace of the similarity transformed representative is therefore the same as the trace of the original representative.

Proof that the characters of two symmetry operations in the same class are identical

The formal requirement for two symmetry operations g and g' to be in the same class is that there must be some symmetry operation f of the group such that $g' = f^{-1}gf$ (the elements g and g' are then said to be *conjugate*). If we consider the characters of g and g' we find:

$$\chi(g') = \text{tr}[\Gamma(g')] = \text{tr}[\Gamma^{-1}(f)\Gamma(g)\Gamma(f)] = \text{tr}[\Gamma(g)\Gamma(f)\Gamma^{-1}(f)] = \text{tr}[\Gamma(g)] = \chi(g) \quad (29.2)$$

The characters of g and g' are identical.

Proof of the Variation Theorem

The variation theorem states that given a system with a Hamiltonian H , then if ϕ is any normalized, well-behaved function that satisfies the boundary conditions of the Hamiltonian, then

$$\langle \phi | H | \phi \rangle \geq E_0 \quad (29.3)$$

where E_0 is the true value of the lowest energy eigenvalue of H . This principle allows us to calculate an upper bound for the ground state energy by finding the trial wavefunction ϕ for which the integral is minimized (hence the name; trial wavefunctions are varied until the optimum solution is found). Let us first verify that the variational principle is indeed correct.

We first define an integral

$$\begin{aligned} I &= \langle \phi | -E_0 | \phi \rangle \\ &= \langle \phi | H | \phi \rangle - \langle \phi | E_0 | \phi \rangle \\ &= \langle \phi | H | \phi \rangle - E_0 \langle \phi | \phi \rangle \\ &= \langle \phi | H | \phi \rangle - E_0 \quad \text{since } \phi \text{ is normalized} \end{aligned} \quad (29.4)$$

If we can prove that $I \geq 0$ then we have proved the variation theorem.

Let Ψ_i and E_i be the true eigenfunctions and eigenvalues of H , so $H\Psi_i = E_i\Psi_i$. Since the eigenfunctions Ψ_i form a complete basis set for the space spanned by H , we can expand any wavefunction ϕ in terms of the Ψ_i (so long as ϕ satisfies the same boundary conditions as Ψ_i).

$$\phi = \sum_k a_k \Psi_k \quad (29.5)$$

Substituting this function into our integral I gives

$$\begin{aligned} I &= \left\langle \sum_k a_k \Psi_k \left| H - E_0 \right| \sum_j a_j \Psi_j \right\rangle \\ &= \left\langle \sum_k a_k \Psi_k \left| \sum_j (H - E_0) a_j \Psi_j \right. \right\rangle \end{aligned} \quad (29.6)$$

If we now use $H\Psi + E\Psi$, we obtain

$$\begin{aligned} I &= \left\langle \sum_k a_k \Psi_k \left| \sum_j a_j (E_j - E_0) \Psi_j \right. \right\rangle \\ &= \sum_k \sum_j a_k^* a_j (E_j - E_0) \langle \Psi_k | \Psi_j \rangle \\ &= \sum_k \sum_j a_k^* a_j (E_j - E_0) \delta_{jk} \end{aligned} \quad (29.7)$$

We now perform the sum over j , losing all terms except the $j = k$ term, to give

$$\begin{aligned} I &= \sum_k a_k^* a_k (E_k - E_0) \\ &= \sum_k |a_k|^2 (E_k - E_0) \end{aligned} \quad (29.8)$$

Since E_0 is the lowest eigenvalue, $E_k - E_0$ must be positive, as must $|a_k|^2$. This means that all terms in the sum are non-negative and $I \geq 0$ as required.

For wavefunctions that are not normalized, the variational integral becomes:

$$\frac{\langle \phi | H | \phi \rangle}{\langle \phi | \phi \rangle} \geq E_0 \quad (29.9)$$

Derivation of the secular equations – the general case of the linear variation method

In the study of molecules, the variation principle is often used to determine the coefficients in a *linear variation function*, a linear combination of n linearly independent functions (f_1, f_2, \dots, f_n) (often atomic orbitals) that satisfy the boundary conditions of the problem. i.e. $\phi = \sum_i c_i f_i$. The coefficients c_i are parameters to be determined by minimizing the variational integral. In this case, we have:

$$\begin{aligned} \langle \phi | H | \phi \rangle &= \langle \sum_i c_i f_i | H | \sum_j c_j f_j \rangle \\ &= \sum_i \sum_j c_i^* c_j \langle f_i | H | f_j \rangle \\ &= \sum_i \sum_j c_i^* c_j H_{ij} \end{aligned} \quad (29.10)$$

where H_{ij} is the Hamiltonian matrix element.

$$\begin{aligned} \langle \phi | \phi \rangle &= \langle \sum_i c_i f_i | \sum_j c_j f_j \rangle \\ &= \sum_i \sum_j c_i^* c_j \langle f_i | f_j \rangle \\ &= \sum_i \sum_j c_i^* c_j S_{ij} \end{aligned} \quad (29.11)$$

where S_{ij} is the overlap matrix element.

The variational energy is therefore

$$E = \frac{\sum_i \sum_j c_i^* c_j H_{ij}}{\sum_i \sum_j c_i^* c_j S_{ij}} \quad (29.12)$$

which rearranges to give

$$E \sum_i \sum_j c_i^* c_j S_{ij} = \sum_i \sum_j c_i^* c_j H_{ij} \quad (29.13)$$

We want to minimize the energy with respect to the linear coefficients c_i , requiring that $\frac{\partial E}{\partial c_i}$ for all i . Differentiating both sides of the above expression gives,

$$\frac{\partial E}{\partial c_k} \sum_i \sum_j c_i^* c_j S_{ij} + E \sum_i \sum_j \left[\frac{\partial c_i^*}{\partial c_k} c_j + \frac{\partial c_j}{\partial c_k} c_i^* \right] S_{ij} + \sum_i \sum_j \left[\frac{\partial c_i^*}{\partial c_k} c_j + \frac{\partial c_j}{\partial c_k} c_i^* \right] H_{ij} \quad (29.14)$$

Since $\frac{\partial c_i^*}{\partial c_k} = \delta_{ik}$ and $S_{ij} = S_{ji}$, $H_{ij} = H_{ji}$, we have

$$\frac{\partial E}{\partial c_k} \sum_i \sum_j c_i^* c_j S_{ij} + 2E \sum_i S_{ik} = 2 \sum_i c_i H_{ik} \quad (29.15)$$

When $\frac{\partial E}{\partial c_k} = 0$, this gives

$$\boxed{\sum_i c_i (H_{ik} - E S_{ik}) = 0} \quad \text{for all } k \quad \text{SECULAR EQUATIONS} \quad (29.16)$$

This page titled [1.29: Appendix A](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

1.30: Appendix B- Point Groups

Non axial groups

$$\begin{array}{c|c} C_1 & E \\ \hline A_1 & 1 \end{array} \quad (1.30.1)$$

$$\begin{array}{c|c|c|c} C_s & E & \sigma_h & \\ \hline A & 1 & 1 & x, y, R_z \\ A' & 1 & -1 & z, R_x, R_y \end{array} \quad \begin{array}{c} x^2, y^2, z^2, xy \\ yz, xz \end{array} \quad (1.30.2)$$

$$\begin{array}{c|c|c|c} C_i & E & i & \\ \hline A_g & 1 & 1 & R_x, R_y, R_z \\ A_u & 1 & -1 & x, y, z \end{array} \quad \begin{array}{c} x^2, y^2, z^2, xy, xz, yz \\ x, y, z \end{array} \quad (1.30.3)$$

C_n groups

$$\begin{array}{c|c|c|c} C_2 & E & C_2 & \\ \hline A & 1 & 1 & z, R_z \\ B & 1 & -1 & x, y, R_x, R_y \end{array} \quad \begin{array}{c} x^2, y^2, z^2, xy \\ yz, xz \end{array} \quad (1.30.4)$$

$$\begin{array}{c|c|c|c|c} C_3 & E & C_3 & C_3^2 & \\ \hline A & 1 & 1 & 1 & x, R_z \\ E & \begin{Bmatrix} 1 & c & c^* \\ 1 & c^* & c \end{Bmatrix} & & & x, y, R_x, R_y \end{array} \quad \begin{array}{c} c = e^{2\pi/3} \\ x^2 + y^2, z^2 \\ x^2 - y^2, xy, xz, yz \end{array} \quad (1.30.5)$$

$$\begin{array}{c|c|c|c|c|c} C_4 & E & C_4 & C_2 & C_4^3 & \\ \hline A & 1 & 1 & 1 & 1 & z, R_z \\ B & 1 & -1 & 1 & -1 & \\ E & \begin{Bmatrix} 1 & i & -1 & -i \\ 1 & -i & -1 & i \end{Bmatrix} & & & & x, y, R_x, R_y \end{array} \quad \begin{array}{c} x^2 + y^2, z^2 \\ x^2 - y^2, xy \\ yz, xz \end{array} \quad (1.30.6)$$

C_{nv} groups

$$\begin{array}{c|c|c|c|c|c} C_{2v} & E & C_2 & \sigma_v(xz) & \sigma'_v(yz) & \\ \hline A_1 & 1 & 1 & 1 & 1 & z \\ A_2 & 1 & 1 & -1 & -1 & R_z \\ B_1 & 1 & -1 & 1 & -1 & x, R_y \\ B_2 & 1 & -1 & -1 & 1 & y, R_x \end{array} \quad \begin{array}{c} x^2, y^2, z^2 \\ xy \\ xz \\ yz \end{array} \quad (1.30.7)$$

$$\begin{array}{c|c|c|c|c} C_{3v} & E & 2C_3 & 3\sigma_v & \\ \hline A_1 & 1 & 1 & 1 & z \\ A_2 & 1 & 1 & -1 & R_z \\ E & 2 & -1 & 0 & x, y, R_x, R_y \end{array} \quad \begin{array}{c} x^2 + y^2, z^2 \\ x^2 - y^2, xy, xz, yz \end{array} \quad (1.30.8)$$

C_{nh} groups

C_{2h}	E	C_2	i	σ_h		
A_g	1	1	1	1	R_z	x^2, y^2, z^2, xy
B_g	1	-1	1	-1	R_x, R_y	xz, yz
A_u	1	1	-1	-1	z	
B_u	1	-1	-1	1	x, y	

(1.30.9)

C_{3h}	E	C_3	C_3^2	σ_h	S_3	S_3^5		$c = e^{2\pi/3}$
A	1	1	1	1	1	1	R_z	$x^2 + y^2, z^2$
E	$\left\{ \begin{array}{cccccc} 1 & c & c^* & 1 & c & c^* \\ 1 & c^* & c & 1 & c^* & c \end{array} \right\}$						x, y	$x^2 - y^2, xy$
A'	1	1	1	-1	-1	-1	z	
E'	$\left\{ \begin{array}{cccccc} 1 & c & c^* & -1 & -c & -c^* \\ 1 & c^* & c & -1 & -c^* & -c \end{array} \right\}$						R_x, R_y	xz, yz

(1.30.10)

D_n groups

D_2	E	$C_2(z)$	$C_2(y)$	$C_2(x)$		
A	1	1	1	1		x^2, y^2, z^2
B_1	1	1	-1	-1	z, R_z	xy
B_2	1	-1	1	-1	y, R_y	xz
B_3	1	-1	-1	1	x, R_x	yz

(1.30.11)

D_3	E	$2C_3$	$3C_2$		
A_1	1	1	1		$x^2 + y^2, z^2$
A_2	1	1	-1	z, R_z	
E	2	-1	0	x, y, R_x, R_y	$x^2 - y^2, xy, xz, yz$

(1.30.12)

D_{nh} groups

D_{2h}	E	$C_2(z)$	$C_2(y)$	$C_2(x)$	i	σ_{xy}	$\sigma(xz)$	$\sigma(yz)$		
A_g	1	1	1	1	1	1	1	1		x^2, y^2, z^2
B_{1g}	1	1	-1	-1	1	1	-1	-1	R_z	xy
B_{2g}	1	-1	1	-1	1	-1	1	-1	R_y	xz
B_{3g}	1	-1	-1	1	1	-1	-1	1	R_x	yz
A_u	1	1	1	1	-1	-1	-1	-1		
B_{1u}	1	1	-1	-1	-1	-1	1	1	z	
B_{2u}	1	-1	1	-1	-1	1	-1	1	y	
B_{3u}	1	-1	-1	1	-1	1	1	-1	x	

(1.30.13)

D_{nd} groups

D_{2d}	E	$2S_4$	C_2	$2C_2'$	$2\sigma_d$		
A_1	1	1	1	1	1		$x^2 + y^2, z^2$
A_2	1	1	1	-1	-1	R_z	
B_1	1	-1	1	1	-1		$x^2 - y^2$
B_2	1	-1	1	-1	1	z	xy
E	2	0	-2	0	0	x, y, R_x, R_y	xy, yz

(1.30.14)

D_{3d}	E	$2C_3$	$3C_2$	i	$2S_6$	$3\sigma_d$		
A_{1g}	1	1	1	1	1	1		$x^2 + y^2, z^2$
A_{2g}	1	1	-1	1	1	-1	R_z	
E_g	2	-1	0	2	-1	0	R_x, R_y	$x^2 - y^2, xy, xz, yz$
A_{1u}	1	1	1	-1	-1	-1		
A_{2u}	1	1	-1	-1	-1	1	z	
E_u	2	-1	0	-2	1	0	x, y	

(1.30.15)

$C_{\infty v}$ and $D_{\infty h}$

$D_{\infty h}$	E	$2C_{\infty}^{\Phi}$...	$\infty\sigma_v$	i	$2S_{\infty}^{\Phi}$...	∞C_2		
Σ_g^+	1	1	...	1	1	1	...	1		$x^2 + y^2, z^2$
Σ_g^-	1	1	...	-1	1	1	...	-1	R_z	
Π_g	2	$2\cos\Phi$...	0	2	$-2\cos\Phi$...	0	R_x, R_y	xz, yz
Δ_g	2	$2\cos 2\Phi$...	0	2	$2\cos 2\Phi$...	0		$x^2 - y^2, xy$
...		
Σ_u^+	1	1	...	1	-1	-1	...	-1	z	
Σ_u^-	1	1	...	-1	-1	-1	...	1		
Π_u	2	$2\cos\Phi$...	0	-2	$2\cos\Phi$...	0	x, y	
Δ_u	2	$2\cos 2\Phi$...	0	-2	$-2\cos 2\Phi$...	0		
...		

(1.30.16)

S_n groups

S_4	E	S_4	C_2	S_4^3		
A	1	1	1	1	R_z	$x^2 + y^2, z^2$
B	1	-1	1	-1	z	$x^2 - y^2, xy$
E	$\begin{Bmatrix} 1 & i & -1 & -i \\ 1 & -i & -1 & i \end{Bmatrix}$				x, y, R_x, R_y	xz, yz

(1.30.17)

S_6	E	C_3	C_3^2	i	S_6^5	S_6		$c = e^{2\pi/3}$
A_g	1	1	1	1	1	1	R_z	$x^2 + y^2, z^2$
E_g	$\begin{Bmatrix} 1 & c & c^* & 1 & c & c^* \\ 1 & c^* & c & 1 & c^* & c \end{Bmatrix}$						R_x, R_y	$x^2 - y^2, xy, xz, yz$
A_u	1	1	1	-1	-1	-1	z	
E_u	$\begin{Bmatrix} 1 & c & c^* & -1 & -c & -c^* \\ 1 & c^* & c & -1 & -c^* & -c \end{Bmatrix}$						x, y	

(1.30.18)

Cubic groups

T	E	$4C_3$	$4C_3^2$	$3C_2$		$c = e^{2\pi/3}$
A	1	1	1	1		$x^2 + y^2, z^2$
E	$\begin{Bmatrix} 1 & c & c^* & 1 \\ 1 & c^* & c & 1 \end{Bmatrix}$					$2z^2 - x^2 - y^2, x^2 - y^2$
T	3	0	0	-1	R_x, R_y, R_z, x, y, z	xy, xz, yz

(1.30.19)

T_d	E	$8C_3$	$3C_2$	$6S_4$	$6\sigma_d$		
A_1	1	1	1	1	1		$x^2 + y^2, z^2$
A_2	1	1	1	-1	-1		
E	2	-1	2	0	0		$2z^2 - x^2 - y^2, x^2 - y^2$
T_1	3	0	-1	1	-1	R_x, R_y, R_z	
T_2	3	0	-1	-1	1	x, y, z	xy, xz, yz

(1.30.20)

Direct product tables

For the point groups O and T_d (and O_h)

	A_1	A_2	E	T_1	T_2
A_1	A_1	A_2	E	T_1	T_2
A_2		A_1	E	T_2	T_1
E			$A_1 + A_2 + E$	$T_1 + T_2$	$T_1 + T_2$
T_1				$A_1 + E + T_1 + T_2$	$A_2 + E + T_1 + T_2$
T_2					$A_1 + E + T_1 + T_2$

(1.30.21)

For the point groups D_4 , C_{4v} , D_{2d} (and $D_{4h} = D_4 \otimes C_i$)

	A_1	A_2	B_1	B_2	E
A_1	A_1	A_2	B_1	B_2	E
A_2		A_1	B_2	B_1	E
B_1			A_1	A_2	E
B_2				A_1	E
E					$A_1 + A_2 + B_1 + B_2$

(1.30.22)

For the point groups D_3 and C_{3v}

	A_1	A_2	E
A_1	A_1	A_2	E
A_2		A_1	E
E			$A_1 + A_2 + E$

(1.30.23)

For the point groups D_6 , C_{6v} and D_{3h}^*

	A_1	A_2	B_1	B_2	E_1	E_2
A_1	A_1	A_2	B_1	B_2	E_1	E_2
A_2		A_1	B_2	B_1	E_1	E_2
B_1			A_1	A_2	E_2	E_1
B_2				A_1	E_2	E_1
E_1					$A_1 + A_2 + E_2$	$B_1 + B_2 + E_1$
E_2						$A_1 + A_2 + E_2$

(1.30.24)

* in D_{3h} make the following changes in the above table

In table	In D_{3h}
A_1	A'_1
A_2	A'_2
B_1	A''_1
B_2	A''_2
E_1	E''
E_2	E'

(1.30.25)

This page titled [1.30: Appendix B- Point Groups](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Claire Vallance](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform; a detailed edit history is available upon request.

Index

A

Abelian

[1.5: Combining Symmetry Operations - 'Group Multiplication'](#)

B

block diagonal

[1.12: Reduction of Representations I](#)

C

character

[1.11: Properties of Matrix Representations](#)

character table

[1.14: Character Tables](#)

chirality

[1.4: Symmetry and Physical Properties](#)

D

degrees of freedom

[1.24: Molecular Vibrations](#)

direct sum

[1.12: Reduction of Representations I](#)

E

electric dipole operator

[1.27: Spectroscopy - Interaction of Atoms and Molecules with Light](#)

G

Great Orthogonality Theorem

[1.15: Reduction of representations II](#)

group multiplication

[1.5: Combining Symmetry Operations - 'Group Multiplication'](#)

L

Little Orthogonality Theorem

[1.15: Reduction of representations II](#)

M

matrix representative

[1.23: A more complicated bonding example](#)

molecular orbital

[1.26: Group theory and Molecular Electronic States](#)

P

point groups

[1.3: Symmetry Classification of Molecules- Point Groups](#)

polarity

[1.4: Symmetry and Physical Properties](#)

projection operator

[1.16: Symmetry Adapted Linear Combinations \(SALCs\)](#)

S

SALC

[1.12: Reduction of Representations I](#)

[1.16: Symmetry Adapted Linear Combinations \(SALCs\)](#)

Schoenflies

[1.3: Symmetry Classification of Molecules- Point Groups](#)

secular equations

[1.21: Solving the Secular Equations](#)

similarity transform

[1.11: Properties of Matrix Representations](#)

symmetry adapted linear combination

[1.12: Reduction of Representations I](#)

[1.16: Symmetry Adapted Linear Combinations \(SALCs\)](#)

symmetry elements

[1.2: Symmetry Operations and Symmetry Elements](#)

symmetry operations

[1.2: Symmetry Operations and Symmetry Elements](#)

U

unfaithful representation

[1.12: Reduction of Representations I](#)

Glossary

Sample Word 1 | Sample Definition 1

Detailed Licensing

Overview

Title: [Symmetry \(Vallance\)](#)

Webpages: 42

All licenses found:

- [CC BY 4.0](#): 76.2% (32 pages)
- [Undeclared](#): 23.8% (10 pages)

By Page

- [Symmetry \(Vallance\) - CC BY 4.0](#)
 - [Front Matter - Undeclared](#)
 - [TitlePage - Undeclared](#)
 - [InfoPage - Undeclared](#)
 - [Table of Contents - Undeclared](#)
 - [Licensing - Undeclared](#)
 - [1: Chapters - CC BY 4.0](#)
 - [1.1: Introduction to Symmetry - CC BY 4.0](#)
 - [1.2: Symmetry Operations and Symmetry Elements - CC BY 4.0](#)
 - [1.3: Symmetry Classification of Molecules- Point Groups - CC BY 4.0](#)
 - [1.4: Symmetry and Physical Properties - CC BY 4.0](#)
 - [1.5: Combining Symmetry Operations - 'Group Multiplication' - CC BY 4.0](#)
 - [1.6: Constructing higher groups from simpler groups - CC BY 4.0](#)
 - [1.7: Mathematical Definition of a Group - CC BY 4.0](#)
 - [1.8: Review of Matrices - CC BY 4.0](#)
 - [1.9: Transformation matrices - CC BY 4.0](#)
 - [1.10: Matrix Representations of Groups - CC BY 4.0](#)
 - [1.11: Properties of Matrix Representations - CC BY 4.0](#)
 - [1.12: Reduction of Representations I - CC BY 4.0](#)
 - [1.13: Irreducible representations and symmetry species - CC BY 4.0](#)
 - [1.14: Character Tables - CC BY 4.0](#)
 - [1.15: Reduction of representations II - CC BY 4.0](#)
 - [1.16: Symmetry Adapted Linear Combinations \(SALCs\) - CC BY 4.0](#)
 - [1.17: Determining whether an Integral can be Non-zero - CC BY 4.0](#)
 - [1.18: Bonding in Diatomics - CC BY 4.0](#)
 - [1.19: Bonding in Polyatomics- Constructing Molecular Orbitals from SALCs - CC BY 4.0](#)
 - [1.20: Calculating Orbital Energies and Expansion Coefficients - CC BY 4.0](#)
 - [1.21: Solving the Secular Equations - CC BY 4.0](#)
 - [1.22: Summary of the Steps Involved in Constructing Molecular Orbitals - CC BY 4.0](#)
 - [1.23: A more complicated bonding example - CC BY 4.0](#)
 - [1.24: Molecular Vibrations - CC BY 4.0](#)
 - [1.25: Summary of applying group theory to molecular motions - CC BY 4.0](#)
 - [1.26: Group theory and Molecular Electronic States - CC BY 4.0](#)
 - [1.27: Spectroscopy - Interaction of Atoms and Molecules with Light - CC BY 4.0](#)
 - [1.28: Summary - CC BY 4.0](#)
 - [1.29: Appendix A - CC BY 4.0](#)
 - [1.30: Appendix B- Point Groups - CC BY 4.0](#)
 - [Back Matter - Undeclared](#)
 - [Index - Undeclared](#)
 - [Glossary - Undeclared](#)
 - [Glossary - Undeclared](#)
 - [Detailed Licensing - Undeclared](#)